

# Index number theory and construction of CPI for complete micro data

Yrjö Vartia  
University of Helsinki, Finland

Antti Suoperä  
Statistics Finland, Helsinki<sup>1</sup>

## Sisällys

Index number theory and construction of CPI for complete micro data .....	1
1. Introduction .....	1
2. Complete micro data and its transformations .....	3
3. Basic concepts and notation for index numbers .....	4
Table 1: Modifications of index numbers. Geometric mean as an example.....	5
4. Laspeyres, Paasche and Palgrave .....	6
5. Three base and observation period weighted means of price relatives.....	7
6. Measuring the <i>Shift</i> of an index number P in respect to Log-Laspeyres .....	14
7. Measuring the <i>Shift</i> of an index number P in respect to Fisher.....	15
8. Simple derivation of the three basic superlative index numbers .....	20
9. Index number production for similar complete micro data .....	22
10. Conclusions .....	23
References .....	25
Appendix 1: Theorems on biasedness and unbiasedness .....	26
Appendix 2: A 10-fork of six basic formulas and their FA´s. ....	31
Appendix 3: TA and FA for the basic six formulas.....	31
Table 1: Time Antithesis formulas TA for the basic six formulas.....	31
Table 2: Factor Antithesis formulas FA for the basic six formulas.....	32

## 1. Introduction

We shall use index number theory only to the extent that helps the construction of CPI in statistical offices, while new complete micro data-sets emerge for commodities. We mean by the complete micro data new data-sets, where both prices and quantities are registered for all homogenous commodities in some market and for all periods. This is often referred as scanner data and is part of Big Data. Complete micro data is different to ‘Billion Prices Project’ used in his CPI by Cavallo and Rigobon (2016) of MIT, because their data is based on vast number of web-prices only (no quantity information). Inclusion of complete micro data in CPI’s will greatly change and simplify the current practices based on Laspeyres formula and complicated rules for elementary aggregates. When the number of commodities increases 50-fold and also the quantities are observed, the index number calculations must change accordingly.

---

<sup>1</sup> We thank for Eugen Koev, Heikki Pursiainen and Timo Koskimäki for their comments and constructive criticism.

The price and quantity indices are the macro equivalents or aggregations of the price and quantity relatives. A price index is roughly speaking some average of the price relatives,  $p_i^{t/0} = p_i^t/p_i^0$ , relevant to an economic unit, say for example consumer. Normally a price index is some well-known weighted average, like arithmetic A, geometric G or harmonic H average, and the weights  $w_i$  are value shares based on either base or observation periods. It is well-known that  $A > G > H$  always (“a three-tined fork of index numbers”) when price relatives vary and their relative differences are approximately equal to half of a variance of relative changes in prices. This result is based on properties of moment means and derived carefully in Vartia (1978) and it contained the critique of Fisher (1922). The main mistakes of Fisher were assuming that Laspeyres and Paasche indices were almost equal and that variances of price and quantity log-changes are essentially the same, which are true only accidentally. The corrected theory is presented in the paper in a more readable and general form than in Vartia (1978). This is a joint work of the authors and continues the co-work of University of Helsinki and Statistics Finland related more generally to aggregation of behavioral functions, see Vartia (2008) and Suoperä and Vartia (2011).

Unfortunately, the selection of the index formulas is quite rarely analyzed purely in terms of the descriptive index number theory. That is, the concept of bias of an index number formula has not been examined accurately, not even in our previous papers. We correct this drawback in chapter 8.

In this paper, we analyze most popular index number formulas and we show how their numerical values depend on the index formula. We show that index number formulas can be classified to formulas having upward or downward bias for small changes compared to index formulas, that must be classified as ‘unbiased’ according to the index number theory. This analysis is carried out for small changes both in log prices and log quantities, and it leads to exact and definite concepts there. The concept of bias vs. unbiasedness is developed carefully in chapter 7. Five first chapters consider the index number problem and CPI on a general level. Appendices consider more technical matters.

One may ask, what makes the index number theory important. Our answer is that it tries to solve the Arrow’s Paradox (concerning the incompatibility of intuitive rules of social decisions) in a quantitative way. The quantity index of consumption (calculated by a precision formula) is constructed to measure the welfare based on consumption in a quantitative way, see Vartia-Weymark (1982) and Vartia (1983). Economists agree to a large extent that it succeeds in its goal. Although there does not exist an ideal (or the only correct) way to measure welfare, it can be measured accurately enough in many important situations. If we agree the set of precision formulas, the welfare change is measured with practical accuracy. Not exactly, but almost. The paper Vartia (1980) takes into account also the saving decisions to the consumer behavior.

In our treatment, we are able to correct some misconceptions held since Fisher (1922), which resulted from his faulty generalizations of his price-quantity data. For example, Laspeyres and Paasche cannot be considered generally even roughly equal. However, Laspeyres formula has been taken as a practical solution in CPI’s, because quantity information is available only for the base period (based on consumer surveys). Even more dramatically a new observation is, that so-called factor antithesis formulas of the basic price indices and their rectified versions become useless, if the variance of the changes in log-quantities is many times larger than the similar variance of prices. This is a probable situation in future very detailed “Big Data”. This case was ignored in Fisher’s authoritative view, where variations in log-changes in prices and quantities were essentially equal.

On the other hand, time antithesis and its use in rectifying formulas remains an excellent idea of Fisher. To our knowledge, this criticism towards “the bible of index numbers” is a new contribution.

The mathematical methods, that we are going to use, are based on well-known references, Fisher (1922), Törnqvist (1936), Stuvell (1957, 1989), Diewert (1976,1978), Vartia (1976, 1978, 1983) and Vartia & Vartia (1984), Törnqvist-Vartia-Vartia (1985). We base our index number theory on basic algebra or arithmetic. No questionable assumptions of “economic behavior” say from consumer theory are needed nor used in our derivations. Economic theorization has a tendency of increasing the list of desired properties of indices, which is usually even too long in the beginning. It is well-known and proved several times, that there cannot

be any index numbers satisfying even certain short and intuitively plausible lists of properties or desiderata. This reminds us of the Arrows paradox in macro decision making. For a comprehensive treatment in index number methods within this approach, see Balk (1995). Hansen-Lucas (1983) documents a quantitative comparison of many important index formulas in the foreign trade of Egypt from 1885-1961.

Our target readers are the index number specialist and constructors in the national statistical offices. We concentrate on those properties of indices, which in our opinion are relevant to these specialists in statistical offices, while they are formulating the new emerging methodology of index construction.

## 2. Complete micro data and its transformations

We verify our mathematical results by numerical analysis. The data used, is well edited drugs data (i.e. self-care drugs) uphold by Pharmaceutical Information Centre Ltd, Finland. It is *complete micro data* in the sense that it contains all prices and quantities (aggregated for the whole country here) for all homogeneous commodities and periods, which are months in the paper.

Instead of couple of dozens product varieties typically included into a CPI sub-index, the data we use contains complete price and quantity data of all periods (here all months during 2013-2016) for about 5000 drugs that require prescription by a medical professional and about 500 self-care drugs (so called VNR-commodities). These are packages of drugs, that have a constant quality over time. This means that problems of quality change do not appear in this data set. Because the actual prices paid by the consumer for prescription drugs are somewhat complicated because of the sickness insurance schemes applied in Finland, we will use only the data on self-care drugs here. It will form our *test data*, which we assume to simulate well (as basic data in CPI) other similar forthcoming complete micro data-sets. We do not consider the peculiarities of this drug data in the paper. In it the prices are rationed and the same in all shops. We consider only features, which we consider to generalize to other complete micro data.

We concentrate on results and proposals that are applicable for similar CPI data having up to 100 times larger sets of commodities than in the current national CPI practices. New and disappearing commodities can also be treated in a systematic and simple way and quality corrections are unnecessary, because the micro commodities are homogeneous. In the new practices, complete price and quantity data appears to become available of practically all commodities in the retail market. We believe that using these new data sets and improved index number methodology, reliability and accuracy of CPI production will be raised to a new level.

In the index number calculations, we split the data in two by five different ways according to how the values of the VNR-commodities have changed between base and observation periods. In the first splitting in two, one part of data 10S consists of the VNR-commodities, whose values have increased or declined more than 10 times and another part 10N is the complement of 10S. In the second split in two the limit is 5, in third 3, in fourth 2 and in the last one 1.33 times increase or decline of the values.

We comment here only the calculations concerning 5N (N for Normal commodities) and 5S (S for Special). The subgroup 5S includes commodities having large relative changes in values (up by multiple 5 or down by 1/5) and thus in quantity or/and in prices. On the other hand, in 5N the values of the commodities stay relatively constant. We show, what kind of effects the special parts 10S, 5S (reported here), 3S, 2S, 1.333S of the data cause to index number calculations. In fact, we shall show some dramatic effects in them. They are so surprising, that one needs some time to digest their consequences.

### 3. Basic concepts and notation for index numbers

We represent the basic concepts of index numbers as an easy reference for experts in statistical offices. Our notation for the index number calculations is the following:

Commodities:	$a_1, a_2, \dots, a_n$ are here self-care drugs and their number $n$ is roughly 500.
Time periods:	$t = 0, 1, 2, \dots$ are the compared situations (only two in binary comparisons).
Prices:	$p_i^t$ is the unit price of $a_i$ in period $t$ .
Quantities:	$q_i^t$ is the quantity of $a_i$ in period $t$ .
Values:	$v_i^t = p_i^t q_i^t$ is the value of $a_i$ in period $t$ .
Total value:	$V^t = \sum v_i^t$ is the total value of all the commodities.
Total value ratio:	$V^{t/0} = V^t / V^0$ is the total value ratio from period 0 to $t$ .
Price relatives:	$p_i^{t/0} = p_i^t / p_i^0$ is the price relative of $a_i$ from period 0 to $t$ .
Quantity relatives:	$q_i^{t/0} = q_i^t / q_i^0$ is the quantity relative of $a_i$ from period 0 to $t$ .
Value relatives:	$v_i^{t/0} = v_i^t / v_i^0$ is the value relative of $a_i$ from period 0 to $t$ .
Value shares:	$w_i^t = v_i^t / \sum_i v_i^t$ is the value share of $a_i$ in period $t$ .

This will be the data, on which the index number calculations are based. Corresponding  $n$ -vectors are denoted by the same symbol without commodity sub-index:

$$(1) \quad p^t, q^t, v^t, p^{t/0}, q^{t/0}, v^{t/0}, w^t$$

This looks much easier: prices, quantities, values. etc.! We assume that all prices and quantities are strictly positive (contain no zeros). This implies that all values, price, quantity and value relatives and values shares are also well-defined and strictly positive. Especially both relatives  $p^{t/0} = p_i^t / p_i^0$  and  $p^{0/t} = p_i^0 / p_i^t$  are defined (no division by zero). This assumption will be relaxed when we later discuss new and disappearing commodities.

Moment mean of order  $\alpha$  (any real number) of strictly positive  $x$ -values and strictly positive weights  $v$  is defined as follows:

$$(2) \quad M^{(\alpha)}(x, v) = \begin{cases} (\sum v_i x_i^\alpha / \sum v_i)^{1/\alpha}, & \text{if } \alpha \neq 0 \\ \prod x_i^{v_i / \sum v_j} = G(x, v), & \text{if } \alpha = 0 \end{cases}$$

This will be a crucial part of our theory, how index numbers deviate from each other. The limit of the moment mean when  $\alpha$  approaches zero is the weighted geometric mean  $G(x, v)$ . Moment mean of non-equal arguments  $x$  is a strictly increasing continuous function of parameter  $\alpha$  from  $M^{(-\infty)}(x, v) = \min(x)$  to  $M^{(\infty)}(x, v) = \max(x)$ . For the important small values of  $\alpha$  the relative change of the moment mean satisfies

$$(3) \quad \log \left( \frac{M^{(\alpha)}(x, v)}{M^{(0)}(x, v)} \right) = \frac{\alpha}{2} s^2(\log x, v) + \frac{\alpha^2}{6} m_3(\log x, v) + \dots$$

where the first two terms of the Taylor series (in respect to  $\alpha$ ) contains the  $v$ -weighted variance and third central moment in the  $\log x$ -scale. This expansion is very accurate for small variances in  $\log x$ . The variance term dominates and gives alone for all moderate values of  $x$  and  $v$  and  $\alpha$  the correct magnitude of the change. The mathematics is so elegant, that we derive the relative deviation of the price index  $M^{(\alpha)}(p^{1/0}, w^0)$  from for the price index  $P = M^{(0)}(p^{1/0}, w^0) = \exp(\sum w_i^0 \Delta \log p_i)$  called Log-Laspeyres  $l$  later. Define  $\hat{p}_i = \log(p_i^{1/0} / P) = \Delta \log p_i - \log P$ , the log-deviations from the mean.

$$\begin{aligned}
(4) \quad & \log \left( \frac{M^{(\alpha)}(p^{1/0}, w^0)}{M^{(0)}(p^{1/0}, w^0)} \right) = \log M^{(\alpha)} \left( \frac{p^{1/0}}{P}, w^0 \right) = \log M^{(\alpha)}(e^{\dot{p}}, w^0) = \frac{1}{\alpha} \log \sum w_i^0 e^{\alpha \dot{p}_i} = \\
& = \frac{1}{\alpha} \log \sum w_i^0 \left( 1 + \alpha \dot{p}_i + \frac{1}{2!} (\alpha \dot{p}_i)^2 + \frac{1}{3!} (\alpha \dot{p}_i)^3 + \dots \right) = \\
& = \frac{1}{\alpha} \log \left( \sum w_i^0 + \alpha \sum w_i^0 \dot{p}_i + \frac{\alpha^2}{2!} \sum w_i^0 \dot{p}_i^2 + \frac{\alpha^3}{3!} \sum w_i^0 \dot{p}_i^3 + \dots \right) \\
& = \frac{1}{\alpha} \log \left( 1 + 0 + \frac{\alpha^2}{2} a_2(\dot{p}, w^0) + \frac{\alpha^3}{6} \sum w_i^0 \dot{p}_i^3 + \dots \right) \\
& = \frac{\alpha}{2} a_2(\dot{p}, w^0) + \frac{\alpha^2}{6} a_3(\dot{p}, w^0) + \dots \quad \text{where } \dot{p}_i = \log \frac{p^{1/0}}{P} = \Delta \log p_i - \log P \\
& = \frac{\alpha}{2} m_2(\ddot{p}, w^0) + \frac{\alpha^2}{6} m_3(\ddot{p}, w^0) + \dots \quad \text{where } \ddot{p}_i = \log p^{1/0} = \Delta \log p_i.
\end{aligned}$$

In this derivation, the log-deviations  $\dot{p}_i = \log \frac{p^{1/0}}{P}$  and their zero mean  $\sum w_i^0 \dot{p}_i = w^0 \cdot \dot{p} = 0$  are the essential points. Here  $a_2(\dot{p}, w^0)$  is the second origo-moment and  $m_2(\ddot{p}, w^0)$  the second central moment. Because trivially (but surprisingly, because of the zero mean of  $\dot{p}$ )  $a_2(\dot{p}, w^0) = m_2(\ddot{p}, w^0) = s^2(\ddot{p}, w^0) = s^2(\dot{p}, w^0) = m_2(\dot{p}, w^0)$ , it is possible to name this crucial parameter in several different ways. This is first confusing.

The derivation is closely related to the moment generating function  $M(\dot{p}, t) = E e^{t\dot{p}} = \sum w_i^0 e^{t\dot{p}_i}$  of the log-deviation variable  $\dot{p}$ , considered as a discrete random variable with probabilities  $p = w^0$ . Its logarithm is the cumulant generating function  $K(\dot{p}, t) = \log M(\dot{p}, t) = \log E e^{t\dot{p}} = \log \sum w_i^0 e^{t\dot{p}_i}$ , which appears above in the form  $\frac{1}{\alpha} K(\dot{p}, \alpha) = \log \sum w_i^0 e^{\alpha \dot{p}_i}$  and the coefficients of its Taylor series gives the sequence of cumulants. The first cumulant is the mean, second the variance and third the third central moment, as above. Higher cumulants are combination of central moments.

Arithmetic, geometric and harmonic means are moment means with parameter 1, 0 and -1, respectively. For small changes, their relative differences from the geometric mean equal half of the variance in the log-argument. For simplicity, we ignore here the third moment term in the analysis of index numbers, see however Vartia (1978).

We represent in a condensed form the operations or modifications of any index number formula in the following table. Time Antithesis TA and Fact Antithesis FA are basic concepts in Fisher's (1922) methodology and we shall review them critically in our paper.

Table 1: Modifications of index numbers. Geometric mean as an example

(1)	(2)	(3)	(4)	(5)
Price Index Number Formula f	f as a Quantity Index Number	TA of f	FA of f	CoF of f
$f_p$	$f_q$	$TA(f_p)$	$FA(f_p)$	$CoF(f)$
$f \begin{pmatrix} p^1 & q^1 \\ p^0 & q^0 \end{pmatrix}$	$f \begin{pmatrix} q^1 & p^1 \\ q^0 & p^0 \end{pmatrix}$	$1/f \begin{pmatrix} p^0 & q^0 \\ p^1 & q^1 \end{pmatrix}$	$V^{1/0}/f \begin{pmatrix} q^1 & p^1 \\ q^0 & p^0 \end{pmatrix}$	$V^{1/0}/f \begin{pmatrix} p^1 & q^1 \\ p^0 & q^0 \end{pmatrix}$
$l_p = \prod \prod \left( \frac{p_i^1}{p_i^0} \right)^{w_i^0}$	$l_q = \prod \prod \left( \frac{q_i^1}{q_i^0} \right)^{w_i^0}$	$p_p = \prod \prod \left( \frac{p_i^1}{p_i^0} \right)^{w_i^1}$	$V^{1/0} / \prod \prod \left( \frac{q_i^1}{q_i^0} \right)^{w_i^0}$	$V^{1/0} / \prod \prod \left( \frac{p_i^1}{p_i^0} \right)^{w_i^0}$

We have presented so-called *Log-Laspeyres* formula  $l_p$  as an example of these modifications. Usually, it is represented in logarithmic form:  $\log l_p = \sum w_i^0 \log \left( \frac{p_i^1}{p_i^0} \right)$ . We just mention for later use two important modifications.

(5) The TA-rectification of  $f = \sqrt{f\left(\frac{p^1 q^1}{p^0 q^0}\right) / f\left(\frac{p^0 q^0}{p^1 q^1}\right)}$  = Cross (geometric mean) with its TA and

(6) the FA-rectification of  $f = \sqrt{f\left(\frac{p^1 q^1}{p^0 q^0}\right)^{V^{1/0}} / f\left(\frac{q^1 p^1}{q^0 p^0}\right)}$  = Cross (geometric mean) ) with its FA.

Starting from  $\log l_p$ , we derive as its TA-rectification the *Törnqvist index*  $\log t_p = \sum \frac{1}{2}(w_i^0 + w_i^1) \log\left(\frac{p_i^1}{p_i^0}\right)$  and as its FA-rectification the index  $\log P = \frac{1}{2}(\sum w_i^0 \log\left(\frac{p_i^1}{p_i^0}\right) + \log \frac{V^1}{V^0} - \sum w_i^0 \log\left(\frac{q_i^1}{q_i^0}\right))$ . We shall show, that the first is unbiased and superlative in a specific sense, but the latter is biased unless the variances of log-changes in prices and quantities happen to be equal. Therefore, Fisher (1922) was in error while considering FA-rectification as a generally valid and effective operation. This is a new result.

#### 4. Laspeyres, Paasche and Palgrave

These are the main building blocks in in number theory. We show their different representations and also try to correct some common misunderstandings. Etienne Laspeyres and Herman Paasche were German economists measuring price changes in Hamburg around 1865 using slightly different methods. Their proposals are the two most important basic indices. As an introduction, we derive for Laspeyres and Paasche price indices their *representations* as weighted averages. For Laspeyres index this is simple:

$$(7) \quad L = L^{1/0} = \frac{p^1 \cdot q^0}{p^0 \cdot q^0} = \frac{\sum p_i^1 q_i^0}{\sum p_i^0 q_i^0} = \frac{\sum \frac{p_i^1}{p_i^0} p_i^0 q_i^0}{\sum v_i^0} = \frac{\sum p_i^{1/0} v_i^0}{\sum v_i^0} = \sum w_i^0 p_i^{1/0}.$$

We have applied *the inner product notation* such as  $p^1 \cdot q^0 = \sum p_i^1 q_i^0$  and  $p^0 \cdot q^0 = \sum p_i^0 q_i^0$ , which is easier to read, because it removes unnecessary summation symbols and product sub-indices. By definition, Laspeyres price index is the cost relative ( $p^1 \cdot q^0$  divided by  $p^0 \cdot q^0$ ) of the old quantity basket  $q^0$ . It can also be calculated as a weighted arithmetic average of the price relatives, where the weights are old values shares. This is called *the practical way of calculating Laspeyres*.

It is applied in national *CPI*'s from the level of its commodity groups by taking  $p_i^{1/0}$  as the value of the price index  $P_i^{1/0}$  of the commodity group and continuing similarly from these. For commodity groups, physical quantities are not available any more, contrary to *elementary aggregates* where they usually exist. This procedure is based on the *consistent aggregation* property Laspeyres formula. Much of common practice in official statistics depends on it, as Pursiainen (2005) stresses.

For the Paasche price index the derivation is similar but more complicated.

$$(8) \quad Pa = Pa^{1/0} = \frac{p^1 \cdot q^1}{p^0 \cdot q^1} = \frac{\sum p_i^1 q_i^1}{\sum p_i^0 q_i^1} = \frac{\sum p_i^1 q_i^1}{\sum \frac{p_i^0}{p_i^1} p_i^1 q_i^1} = \frac{\sum v_i^1}{\sum p_i^{0/1} v_i^1} = \frac{1}{\sum w_i^1 p_i^{0/1}}$$

By definition, Paasche price index is the cost relative ( $p^1 \cdot q^1$  divided by  $p^0 \cdot q^1$ ) of the new quantity basket  $q^1$ . It can also be calculated as a weighted *harmonic* average of the price relatives, where the weights are new values shares. In practice, this is the way it is calculated. It can be used only if the new value shares are known, which is a strong condition. In our complete micro data on drugs, also new quantities are known and new values shares can be calculated. This holds for any scanner-type new micro data and it changes strongly the whole methodology of CPI calculation.

Note also that Laspeyres and Paasche are based on the same basket idea of calculation:  $L$  is based on the old basket of goods  $q^0$ , while  $Pa$  uses the new basket  $q^1$ . Paasche may be defined actually by Laspeyres calculated in the reverse direction  $1 \rightarrow 0$ , i.e. it is evidently the inverse of

$$(9) \quad L^{0/1} = \frac{p^0 \cdot q^1}{p^1 \cdot q^0}.$$

This dependence means actually, that Paasche is the *Time Antithesis* of Laspeyres,  $Pa = TA(L)$ . This concept is introduced in detail in chapter 8.

An elementary but quite common mistake is to imagine, that the Paasche price index as a weighted *arithmetic* average of the price relatives weighted by new values shares. This is not Paasche, but the index number proposed by an economist Palgrave in 1868. It does not have a quantity basket interpretation, like Laspeyres and Paasche. Palgrave has two doses of upward bias compared to Paasche. Only if Paasche is strongly biased down, Palgrave index as such is applicable. As shown by Vartia-Vartia (1978) an average of Laspeyres and Palgrave is *always* biased upwards compared to Fisher (geometric average of Laspeyres and Paasche) or to other unbiased formulas.

This kind of error was made e.g. in the Bank Finland in the choice of the formula in the Bank of Finland Currency Index, see Vartia-Vartia (1978). Economists in the Bank of Finland evidently thought, that they calculated Edgeworth index. It exaggerated the revaluation in Finnish currency by the amount of the bias of the index. The currency index increased more than was the actual case. The bias and claimed unintentional revaluation of Finnish mark had increased from 1974 to 1984 to 2.4%. Concretely, a corresponding devaluation of 2.4% in 1984 would have been needed to eliminate the discrepancy for 1984. This large devaluation (which would have corrected the actual situation to the officially declared one) would have meant an extra income of roughly FIM 1500 million (250 million euros in 1984 prices) only in 1984 to the export sector. Thus, the export sector of the economy was officially claimed to get systematically more foreign currency than they actually got. Official data did not correctly describe the reality. As this example shows, biases and other weaknesses of index calculations are not unimportant details in the macro-economic analysis. The formulas and weights must be chosen correctly in official indices.

## 5. Three base and observation period weighted means of price relatives

Irving Fisher (1922) visualized the index numbers as forks containing a certain number of tines. This is an important visualization of differences between various indices and other choices of calculation. In his data, various index numbers happened to fall into five quite separate groups or “tines”, which formed “the five-tined fork”, a kind of quantum theory of index numbers. Much of this was accidental and resulted because  $L$  and  $P$  happened to be near each other, which is not generally or even usually true. Fisher regarded these formulas as “very good”, which was a mistake as will be shown below. Fisher’s “Five-tined Fork” with doses of bias (or quanta) =  $0, \pm 1, \pm 2$  as a realistic “quantum theory” of index numbers was criticized already in Vartia (1978), but it did not raise much attention. The concepts of bias of an index number was defined for the first time along these lines in Vartia (1978) and Vartia-Vartia (1984) as a small deviation concept, but evidently too loosely. Also, this point was bypassed unnoticed among index number literature. Probably it was not understood, *how* it differed from the Fisher’s concept of bias and unbiasedness. These concepts are now presented here in chapter 7 and in Appendix 1 once again, but more accurately.

Laspeyres has been regarded as the “correct prototype” of all index calculations all over the world both as a price and quantity index because of its simplicity and easy applicability and despite of all sorts of opposing evidence on its upward bias and asymmetry.

Fisher (1922) made some wrong generalizations on these matters. Now almost 100 years later, it is time to build a corrected index number theory on the following foundations: how various choices on averaging and weighting affect index numbers and where the unbiased indices situate. Now the analysis can be based on

modern mathematics (moment means, approximations, function theory, knowledge of new “superlative” or “ideal” index numbers), large data sets and computing power. We must, however, appreciate here Fisher’s genius, as his concepts and classifications still after 100 years govern the development of index theory.

Next, we define three price indices based on arithmetic, geometric and harmonic means of price relatives and old value shares. The three-tined fork having base period weights contains the following indices (three weighted averages  $A$ ,  $G$  and  $H$ ):

$$\begin{aligned}
 (10) \quad L &= A(p^{1/0}, w^0) = \sum w_i^0 p_i^{1/0} = \frac{p^1 \cdot q^0}{p^0 \cdot q^0} && \text{“Laspeyres”} \\
 l &= G(p^{1/0}, w^0) = \prod \left(\frac{p_i^1}{p_i^0}\right)^{w_i^0} = \exp\left(\sum w_i^0 \log p_i^{1/0}\right) && \text{“Log-Laspeyres”} \\
 Lh &= H(p^{1/0}, w^0) = 1/\sum w_i^0 p_i^{0/1} && \text{“Harmonic Laspeyres”}
 \end{aligned}$$

We also use the upper double-dot to denote log-change, as in  $\ddot{p}_i = \Delta \log p_i = \log(p_i^1/p_i^0)$  and similarly for quantities  $\ddot{q}_i$  and values  $\ddot{v}_i$ . Logarithms are natural, of course, see Törnqvist-Vartia-Vartia (1985). Single upper dot is reserved for log-deviations from the mean as in Vartia (1978), e.g.  $\dot{p}_i = \Delta \log p_i - \log P = \dot{p}_i - \log P$ .

More symmetrically written  $\log l = \sum w_i^0 \log(p_i^1/p_i^0) = \sum w_i^0 \ddot{p}_i = \ddot{l}$ . Using (3-4) for moments means, the relative differences of  $L = M^{(1)}(p^{1/0}, w^0)$  and of  $Lh = M^{(-1)}(p^{1/0}, w^0)$  from of  $l = M^{(0)}(p^{1/0}, w^0)$  are approximately half of the “old” variance  $s_0^2(\dot{p}) = s^2(\dot{p}, w^0)$  in the deviations of log-prices<sup>2</sup>. We call  $s_0^2(\dot{p})$  shortly the old variance of  $\dot{p}$ .

We have always  $\log L > \log l > \log Lh$  for non-equal price relatives or  $\dot{p}$ . More accurately, we have  $\log L = a + \Delta_0 > a > \log Lh = a - \Delta_0$ , where  $a = \log l$  and  $\Delta_0 = \frac{1}{2}s_0^2(\dot{p})$ .

The three-tined fork with the observation period weights contains also three weighted averages:

$$\begin{aligned}
 (11) \quad Pl &= A(p^{1/0}, w^1) = \sum w_i^1 p_i^{1/0} && \text{“Palgrave”} \\
 p &= G(p^{1/0}, w^1) = \prod \left(\frac{p_i^1}{p_i^0}\right)^{w_i^1} = \exp\left(\sum w_i^1 \log p_i^{1/0}\right) && \text{“Log-Paasche”} \\
 Pa &= H(p^{1/0}, w^1) = 1/\sum w_i^1 p_i^{0/1} = \frac{p^1 \cdot q^1}{p^0 \cdot q^1} && \text{“Paasche”}
 \end{aligned}$$

More symmetrically written  $\log p = \sum w_i^1 \log(p_i^1/p_i^0) = \sum w_i^1 \ddot{p}_i = \ddot{p}$ . The logarithmic differences of  $Pl$  and  $Pa$  from  $p$  are half of the “new” variance  $s_1^2(\dot{p}) = s^2(\dot{p}, w^1)$  of the changes of log-prices  $\dot{p} = \Delta \log p$ . We call  $s_1^2(\dot{p})$  shortly the new variance of  $\dot{p}$ .

We have always  $\log Pl > \log p > \log Pa$  for non-equal price relatives or  $\dot{p}$ . More accurately, we have  $\log Pl = b + \Delta_1 > b > \log Pa = b - \Delta_1$ , where  $b = \log p$  and  $\Delta_1 = \frac{1}{2}s_1^2(\dot{p})$ .


<sup>2</sup> The same variance may be written also  $s^2(\dot{p}, w^0)$  using the log-changes  $\dot{p}_i = \Delta \log p_i = \log(p_i^1/p_i^0)$ .

<sup>3</sup> Log-Paasche and log-Laspeyres are also called Geometric Paasche and Geometric Laspeyres, especially if confusion to the logarithms of Paasche  $\log Pa$  and Laspeyres  $\log L$  are at stake.



The old and new variances  $s_0^2(\hat{p})$  and  $s_1^2(\hat{p})$  govern the log-differences of the three indices of the old and new forks.

The problem remains how the three-tined forks of the old and new weights situate in respect to each other. As data shows, there does not exist any firm geometric rules. Especially, Fisher's 'Five-tined Fork' was only an accident, not any rule. In our data for the self-care medicines, Laspeyres and Paasche had for every month large biases up and down, respectively. They did not lie in the center but on the opposite sides of the index fork and they were the most biased of these six formulas all the time. Also, the number of tines is not five, as in Fisher's data, but varies between the extremes three and six, see figure below.




As the figure shows, the type of the fork that the six indices form, varies from month to month. But here the Laspeyres-type fork with old weights, Laspeyres on the top, is surprisingly always higher than the new weight fork, where Paasche is the lowest index. All other basic indices remain between  $L$  and  $Pa$ .

Fisher's Five-tined Fork consists of five groups of indices with typical representatives

$Pl > p > Pa \approx L > l > Lh$ . Fisher claimed that  $Pa \approx L$  are unbiased (and very good) index numbers, while all other of these basic indices have  $\pm 1$  or  $\pm 2$  doses of bias. This theory of Fisher's is only one possibility, not a generally valid theory. The assumption  $Pa \approx L$  does not hold generally. For instance, the figure 1 above provides a counter-example, where  $Pa$  and  $L$  are not in the middle tine but the lowest and highest indices. Here the substitution effects are exceptionally large. Fisher's theory of Five-tined Fork is rejected. For more information, google Vartia (1978).

The superlative formulas  $F$ ,  $Stu$ ,  $MV$  and  $t$  form a much tighter band within the six indices here:




All sensible or minimally accurate index numbers should remain within these limits.

As a warning, we show a picture concerning the six factor antithesis formulae, like  $FA(l) = V^{1/0}/l(q)$  and  $FA(p) = V^{1/0}/p(q)$ . This factor antithesis of  $l$  arises by *first* using the same formula for quantities,  $l(q)$ , instead of prices, and *then* returning back to the price space by calculating its co-factor  $CoF(l(q)) = V^{1/0}/l(q) = FA(l)$ . As a technical operation, this is simple, but *conceptually* it requires some expertise in index numbers. Note that the log-difference of two  $FA$ 's depends only on the difference of the formulas in the quantity side, say:  $\log \frac{FA(p)}{FA(l)} = \log \frac{V^{1/0}}{p(q)} - \log \frac{V^{1/0}}{l(q)} = \log \frac{l(q)}{p(q)}$ . Therefore, the  $FA$ 's of the six basic indices vary much more than their original price equivalents. The  $FA$ 's for the sex indices have represented in Appendix two.


The  $FA$ 's of the six basic formulas do not remain within the limits of the picture above and it shows that there is something seriously wrong with them. They are based on applying the six basic indices in the quantity side of the monthly data, where the variance of the quantity changes is roughly 35 times bigger than the similar price variance. Therefore, the  $FA$ 's of the six formulas have index forks dispersing roughly 35 times more than the ordinary price index forks in our data. In Fisher's yearly test data, this was not the case, but instead the variations in price and quantity relatives were exceptionally almost the same. Many of Fisher's conclusions and suggestions based on the Factor Antitheses  $FA$  and on the symmetrical treatment of  $p$ 's and  $q$ 's are based on this accidental and normally false condition.

We cannot get support from the quantity side of data, because it disperses so heavily!



Note how Laspeyres and Paasche appear in the middle of the picture. They are OK and everything else is out of lines.

In the next picture, we show all the six basic indices and their FA's in the figure. Note how much tighter the original six price indices move compared to their FA's.




We will comment later some Fisher's false conclusions which depend on this exceptional feature of his data. In practice, this means that price indices derived from quantity relatives, so called factor antithesis formulas

for prices, become less and less useful as the variance of the changes in *quantities* increases. These effects are no minor nuances, because FA-indices in the sub-group of special commodities 5S will be found literally useless (except  $L$  and  $Pa$ , see Appendix 2).


As a concrete conclusion, the factor antithesis formulas of the basic six price index formulas are as such useless formulas because of their large biases (about 35 times larger than for the basic indices). This rejection of FA formulas does not hold for the FA's of  $L$  and  $P$ , because their FA's are the same indices but in the reverse order:  $FA(L) = P$  and  $FA(P) = L$ . Check that these indices satisfy  $L * P(q) = P * L(q) = \frac{V^1}{V^0}$  with obvious notation. Prove that for Fisher's indices  $F * F(q) = \frac{V^1}{V^0}$ . Hint: multiply the former equations, group their terms and take a square root.

Price indices should not be calculated as functions of the same quantity indices (or cofactors) of the six basic formulas (or their rectifications). Indices calculated via quantity data varied roughly in the same proportion as the variances  $s^2(\dot{p}, w)$  and  $s^2(\dot{q}, w)$  of price and quantity log-changes!


The FA's of the six basic indices (and their derivations, except the ones based on  $L$  and  $Pa$ ) become totally useless in for the group 5S of special commodities.



Also, here  $L$  and  $P$  are OK, and everything else is out of bounds. Factor antithesis indices vary from 0,01 to 1000, as the basic price indices vary within 0,9 - 1,1, as shown below.



The precision formulas are still much more accurate. They do an excellent job also in the set 5S of special commodities having strongly variable values. They vary systematically and their differences are maximally only three percent.




## 6. Measuring the *Shift* of an index number $P$ in respect to Log-Laspeyres

We transform the figure above to a form which better shows the deviations of its all indices from the *Log-Laspeyres*  $l$ . We want to compare indices for different periods and for hypothetical data, where changes in prices and quantities approach zero or some proportional values. Therefore, we divide the relative difference of the compared indices by the average relative difference of our six basic indices. We define the *shift*  $S$  of an index number (say  $p$ ) in respect to *Log-Laspeyres*  $l$  as the difference of  $p$  and  $l$  in the log-scale divided by the average deviation of times from the central time:

$$(12) \quad S(p) = Shift(p, l) = \frac{\log(\frac{p}{l})}{\frac{1}{4}(\log(\frac{L}{Lh}) + \log(\frac{Pl}{P}))} \cong \frac{\log(\frac{p}{l})}{\frac{1}{2}s^2(\dot{p}; \bar{w})}.$$


$Shift(P, l)$  is the relative difference of any index  $P$  from  $l$  divided by the average distance between the six nearby indices of our two forks. These forks consist of 6 different linear approximations for the superlative indices like Fisher or Törnqvist. Here the denominator, half the variance of  $\dot{p}$ , acts as a magnifying glass which expands the picture when the deviations of  $\dot{p}$  are small or, especially, when  $s^2(\dot{p}; \bar{w})$  approaches zero and the 6 basic indices approach the same value. Note that, although  $Shift(P, l)$  approaches the form  $0/0$ , its limit value is usually a small real number.

For the indices  $(L, l, Lh)$  of the old fork the shifts become  $(S(L), S(l), S(Lh)) \cong (1, 0, -1)$  with essential equality at least for small deviations in log-changes of prices and quantities. For the indices  $(Pl, p, P)$  of the new fork the shifts  $(S(Pl), S(p), S(Pa))$  are more interesting. They show, how the new fork deviates (up or down) from the old fork in the normalized scale. For the first period, we have a four-tined fork and for the last a five-tined fork. All the forks differ much from Fisher's data where  $L \approx Pa$  as here  $L$  and  $Pa$  are the largest and lowest figures. We will show how we can get rid of all assumptions related to their sizes and mutual differences.



The small fluctuations in the above figure disappear completely, if we consider the asymptotic case, where the changes in the prices approach zero or some proportional values. These can be calculated by somewhat

complicated algebra, see Appendix 1. The case  $S(p) = Shift(p, l) \cong 0$  where  $l$  and  $p$  happen to be almost equal, looks as follows:



The shift parameter  $S$  of the 6 indices varies from 1 to -3 and the combined fork has 3, 4 or 5 times, as the Paasche-type indices (new 3-fork) shift downwards in this data. This data clearly contradicts Fisher's main conclusion, that  $L$  and  $Pa$  are unbiased and "very good" index numbers. This conclusion was as a general advice a mistake while they may be strongly biased, as in this data. None of our six indices cannot be classified as unbiased, because in some data their biases are clearly realized. But we shall show in a minute that their rectifications based on their time antithesis are all unbiased and superlative index numbers for small price and quantity changes (not necessarily generally).

## 7. Measuring the *Shift* of an index number $P$ in respect to Fisher

One of the basic features of Fisher's theory is to relate all other index numbers to it. This makes sense perfectly for small changes, but can be questioned for large ones. The *Shift* of an index number formula  $P$ , say,  $Shift(P; \cdot)$ , is naturally defined as a shift  $S$  in respect to Fisher, Törnqvist or any other *asymptotically unbiased* index number formula. This helps us to define the *bias* of an index number for small changes or asymptotically. The question of *bias* must be left open for moderate or large changes. Therefore, we call our function symmetrically as a shift between two formulas. To be explicit, we decide to use the traditional Fisher index as the basic point of reference. The expression for  $Shift(P; F)$  for actual or moderate changes is defined as follows

$$(13) \quad Shift(P, F) = \frac{\log\left(\frac{P}{F}\right)}{\frac{1}{4}\left(\log\left(\frac{L}{Lh}\right) + \log\left(\frac{Pl}{P}\right)\right)} \cong \frac{\log\left(\frac{P}{F}\right)}{\frac{1}{4}(s^2(\dot{p}, w^0) + s^2(\dot{p}, w^1))} \cong \frac{\log\left(\frac{P}{F}\right)}{\frac{1}{2}s^2(\dot{p}, \bar{w})}.$$

Note that this is an anti-symmetric function:  $Shift(F, P) = -Shift(P, F)$ . Irrespective of the size of the denominator, some values of  $Shift(P, F)$  for our six basic indices  $P$  must be outside or at the limits of the interval  $(-1, +1)$ . Think of the case  $log l \approx log p$  shown above, where the two three-forks essentially coincide.  $Shift(P, F)$  is well-defined and attains similar values also when its denominator or, more radically, all commodity deviations in log-changes of prices and quantities tend to zero (almost proportional changes APC). These are needed in the definitions of asymptotic bias below.

An index number  $P$  is called *unbiased and superlative for small changes* only if  $Shift(P, F) \rightarrow 0$  or its numerator approaches zero quicker than its denominator. This happens for indices, which are quadratic approximations of  $F$  for small changes in log-prices and log-quantities. Fisher's theory of *Five-tined Fork* was clearly mistaken, see Vartia (1978) and Vartia-Vartia (1984), and no replacement has been generally presented this far. These concepts aim at developing such a new theory.

The function  $Shift(P, F)$  takes the role of magnifying glass when its denominator is small. It proves to be essential how it behaves, when its denominator approaches zero. We classify the index numbers expressed in terms of log-changes  $P = f\left(\frac{p^1}{p^0} \frac{q^1}{q^0}\right) = P(\check{p}, \check{q}, w^0)$  by the way the shift function  $S(\check{p}, \check{q}, w^0) = Shift(P, F)$  behaves for small values of its log-changes  $\check{p}, \check{q}$ . Especially important is, whether this function has a **unique limit value** when these arguments tend to zero or not. This determines whether the price index formula  $P = P(\check{p}, \check{q}, w^0)$  is **unbiased and superlative, permanently biased, contingently biased or freakish for small changes**. We formulate the concept of *bias and unbiasedness* of an index number formula more accurately in two different ways:

An index number  $P$  is said to be *superlative for small changes* (for SC)  
if and only if

its  $Shift(P, F)$  in respect<sup>4</sup> to  $F$  **approaches a unique limit** and it is zero, when all the log-changes in  $p$ 's and  $q$ 's tend to zero. The same zero limit must be attained by all different paths of the inputs or independently how they approach zero.

An index number  $P$  is said to be *permanently biased for small changes* (SC)  
if and only if

its  $Shift(P, F)$  in respect to  $F$  **approaches a unique limit** which is a non-zero real number, when all the log-changes in  $p$ 's and  $q$ 's tend to zero. The same non-zero limit must be attained by all different paths of the inputs. This non-zero value is the value of the permanent bias (a fraction of the norm up or down compared to  $F$ ) in the index.

An index number  $P$  is said to be *contingently biased for small changes* (SC)  
if and only if

its  $Shift(P, F)$  in respect to  $F$  **does not approach a unique limit**, when all the log-changes in  $p$ 's and  $q$ 's tend to zero. Different real numbers are approached for different paths of the inputs approaching zero.

An index number  $P$  is said to be *freakish for small changes* (SC)  
if and only if

its  $Shift(P, F)$  in respect to  $F$  **does not approach a unique limit**, when all the log-changes in  $p$ 's and  $q$ 's tend to zero. Instead the  $Shift(P, F)$  approaches  $\pm\infty$  for some paths of the inputs approaching zero. It is also contingently biased, in a very serious way.

There are only four groups of index numbers:

- 1 **Superlative** with existing  $\lim_{SC} Shift(P, F)$  equaling zero. These are quadratic approximations of  $F$ . They do not differ from each other for small changes.
- 2 **Permanently biased** with existing  $\lim_{SC} Shift(P, F)$  equaling a non-zero real number, say  $b$ . They are permanently biased up or down by  $b$ . These differ from  $F$  as follows:  
 $LogP = logF + b * s^2(\check{p}, \check{q})$  +third and higher order terms.
- 3 **Contingently biased** where  $\lim_{SC} Shift(P, F)$  does not exist, because different real numbers are approached for different paths of  $\check{p}, \check{q}$  approaching zero. For these  $LogP$  is only a linear approximation of  $logF$ , i.e. it differs from  $logF$  by quadratic and higher order terms. They differ from  $F$  in a haphazard way even for small values of  $\check{p}, \check{q}$  depending on the contingent features of the data.
- 4 **Freakish** where  $\lim_{SC} Shift(P, F)$  does not exist, because infinities are approached for some paths of  $\check{p}, \check{q}$  approaching zero. For these  $LogP$  is not even a linear approximation of  $logF$ , but even a worse one. Typically, a freakish index  $P$  is evenly weighted or the

<sup>4</sup> or equivalently in respect to any other superlative formulas such as  $t, D, Dr, SV \dots$



weights are based on a third period or are chosen independently of the old or new value shares, like in Lowe's formula, see Vartia (1976, p 60-61). Also, freakish indices are **contingently biased**, in a very serious way.

There is no reason of using freakish, permanently biased or contingently biased formulas for complete micro data, because there are an infinite number of superlative index numbers, where one can choose of, see Appendix 1. There are no finite steps or quanta by which freakish, permanently biased or contingently biased formulas can only differ from  $F$  or its quadratic approximations. These formulas differ continuously from  $F$ : there does not exist a valid "quantum theoretic" correction of Fisher's mistaken Five-tined Fork. The whole idea of Fisher was based on classifying only proposed or other apparent formulas, not all possible formulas. For instance, any weighted mean of  $l$  and  $L$  (say  $0.1l + 0.9L$ ) is a valid contingently biased formula. Similarly, any weighted mean of  $t$  and  $F$  (say  $0.1t + 0.9F$ ) is a valid superlative formula.

Fisher's Five-tined Fork has been an excellent selling argument, which succeeded to sell Laspeyres contingently biased and suspicious formula  $L$  as a standard formula in public index number production. It took almost a century to show convincingly that it was good selling argument but against the facts.

This was the idea of definition of unbiasedness in Vartia (1978) and Vartia-Vartia (1984), which was formulated partly verbally and did not raise much attention. In vector notation, the smallness condition means that the (unweighted Euclidean) length of vectors  $\|\Delta \log p\| = \sqrt{\sum (\Delta \log p_i)^2} \rightarrow 0$  and  $\|\Delta \log q\| \rightarrow 0$ . This implies e.g.  $\|\Delta \log v\| \rightarrow 0$  and  $\|\Delta \log w\| \rightarrow 0$ . Also  $\log P \rightarrow 0$ ,  $\log Q \rightarrow 0$  and  $\log \frac{v^1}{v^0} \rightarrow 0$ . All these conditions correspond to "small changes".

The second definition of *bias and unbiasedness* is a stricter one and is related to almost proportional changes, not to almost zero changes as above. Here an index number  $P$  must be almost equal (a quadratic approximation) to  $F$  to be unbiased and superlative for APC already when the log-changes of prices and quantities approach any proportional values. This condition of inputs can be described as "the commodity variation in log-prices and log-quantities vanishes" or the variances  $s^2(\dot{p}, \bar{w})$  and  $s^2(\dot{q}, \bar{w})$  tend to zero. Price and quantity indices or their log's  $\log P$  and  $\log Q$  may attain any value, not zero as above.

An index number  $P$  is said to be **superlative for almost proportional changes** (for APC) if and only if

its  $Shift(P, F)$  in respect<sup>5</sup> to  $F$  **approaches a unique limit** and it is zero, when all log-changes of prices and quantities approach any proportional values. The same limit must be attained by all different paths of the inputs. Now the variances  $s^2(\dot{p}, \bar{w})$  and  $s^2(\dot{q}, \bar{w})$  of the log-changes in  $p$ 's and  $q$ 's around their mean values both tend to zero, while  $\log P$  and  $\log Q$  may get any constant values.

An index number  $P$  is said to be **permanently biased for almost proportional changes** (APC) if and only if

its  $Shift(P, F)$  in respect to  $F$  **approaches a unique limit** which is a non-zero real number, when all log-changes of prices and quantities approach any proportional values. The same non-zero limit must be attained by all different paths of the inputs. This non-zero value is the value of the permanent bias (a fraction of the norm up or down compared to  $F$ ) in the index.

An index number  $P$  is said to be **contingently biased for almost proportional** (APC) if and only if

its  $Shift(P, F)$  in respect to  $F$  **does not approach a unique limit**, when all log-changes of prices and quantities approach any proportional values. Different real numbers are approached for different paths of the inputs approaching constants.

<sup>5</sup> or equivalently in respect to any other asymptotically unbiased or superlative formulas such as  $t$ ,  $D$ ,  $Dr$ ,  $SV$  ...

An index number  $P$  is said to be *freakish for almost proportional changes (APC)* if and only if


its  $Shift(P, F)$  in respect to  $F$  **does not approach a unique limit**, when all log-changes of prices and quantities approach any proportional values. The  $Shift(P, F)$  approaches  $\pm\infty$  for some paths of the inputs approaching constants. It is also contingently biased, in a very serious way.

The second definition is stronger and demands more and is presented here only for later reference and in order to give some clarifying comments. If an index is superlative for almost proportional changes, it is necessarily superlative for small changes. It is rather remarkable, that e.g.  $F$  and  $t$  approximate each other equally well irrespective of how much the prices and quantities have changed (i.e. how large their proportional values have been).  $MV$  are superlative for  $SC$ , but it is not superlative for  $APC$ .


We use mostly to the first definition, which seems to conform best to the current research in index number theory.

In Appendix 1 one we formulate 14 theorems and their proofs. By these theorems we may classify the index number formulas in to superlative indices, permanently, contingently biased and freakish indices. They also verify our classification of index numbers in to four groups.

Shifts and biases get large values if also  $FA$ 's of our basic formulas are taken into account.



Especially  $FA(Lh)$  and  $FA(PI)$  contain huge biases. If  $FA$ 's are dropped, the shifts and biases near the zero-line above become visual:




These figures show, that

- A. the index forks are not stationary in time and
- B. Fisher's Five-tined Fork does not appear at all here, not to speak as the only possibility.

Fisher's theory is clearly rejected.

Inside these bounds, we find all the superlative formulas, which have zero biases for SC although their shifts from  $F$  differ slightly from zero (because changes are not infinitesimal):



We stress that the normed deviation from Fisher, the limit of  $Shift(P, F)$ , either up or down is a clear fault of formula  $P$  if the changes in prices and quantities are small.

The shift formulas are related to the same denominator norm, which is calculated from four extreme members of the forks ( $l$  and  $p$  are not needed). It is a good practice always to calculate all the six basic indices (not just  $L$  or perhaps  $L$  and  $P$ ). This allows calculating the shift and the many superlative indices like Törnqvist (and its derivatives), Fisher, Drobish, Stuel et al. from them.

The two three-forks  $(L, l, Lh)$  and  $(Pl, p, Pa)$  provide a natural basis for all *quantitative* comparisons (such as  $Shift(p, l)$  and  $Shift(P, F)$ ) of index number formulas. Such data would increase the applications of official data, perhaps to surprising directions.

Fisher thought that biases of formulas lie within limits of  $(-2, +2)$ . This is false. As shown above, biases of some FA-formulas can be large real numbers, less than -300 or more than 300.

## 8. Simple derivation of the three basic superlative index numbers

Table 1 in Appendix 3 shows that the TA formulas of the original formulas remain in the same set, but their order is reversed. This is essential: For instance, the highest index of the old fork, i.e.  $L$ , is mapped to the lowest index of the new fork or  $Pa$ . It is easier to grasp this algebraic structure in terms of geometry or in the picture, where both forks are presented. In our opinion, this is perhaps the best idea of Fisher. Especially its use in rectification of the formulas gives superlative results!

We will show, that the rectified formulas (in respect to its time antithesis TA) will lead to the same value for small variances of  $\dot{p}$  and  $\dot{q}$ . This point served as the main point in the definition of *asymptotic unbiasedness* of index numbers, which is an exact concept for small *deviations* in  $\dot{p}$  and  $\dot{q}$ , i.e. how much these log-changes *differ from some constants*. Their *magnitude* (i.e. the values of the price and quantity indices or constants) is totally irrelevant! For large deviations of  $\dot{p}$  or large deviations of  $\dot{q}$  (or both) the question of unbiasedness is left slightly ambiguous. This main message of Vartia (1978) and Vartia-Vartia (1984) was left unnoticed in index number literature. But it still clarifies many problems in index numbers.

The whole idea is the same as in feedback in dynamic systems (see Appendix 2, Table 1): a formula  $(1) = (2)$  is used first in the other direction  $(3)$ , and then returned back to the original comparison  $(4) = (5)$ . This is used as the correcting information for the original formula  $(1) = (2)$ . This is the feedback connection used in *rectifying by the time antithesis*. We will show that *TA-rectified formula*

$$(14) \quad \sqrt{f \begin{pmatrix} p^1 & q^1 \\ p^0 & q^0 \end{pmatrix} / f \begin{pmatrix} p^0 & q^0 \\ p^1 & q^1 \end{pmatrix}}$$

gives almost the same results (for small *deviations* in  $\dot{p}$  = the log-changes of prices and  $\dot{q}$  = the log-changes of quantities) for all our basic indices. This 'constant mean point' serves as *the reference* for all formulas that are called *asymptotically unbiased*, as defined in chapter 6.

For small deviations in  $\dot{p}$  and  $\dot{q}$  or regular moderate deviations, the old and new forks have (approximately) the same dispersions (= deviations or separations of the three tines) in the log-scale, as stated by Theorems 1-3 (see Appendix 1). We describe this asymptotic situation of small changes below and derive simply its main conclusions.

In the base fork, the three log-indices differ approximately by half the old price variance  $\Delta^0 = \frac{1}{2}S_0^2(\dot{p})$ :

$$\begin{array}{ll} \log L = a^0 + \Delta^0 & TA(L) = Pa \\ \log l = a^0 & TA(l) = p \\ \log Lh = a^0 - \Delta^0 & TA(Lh) = Pl \end{array}$$

In the new fork, the three log-indices differ approximately by half the new price variance  $\Delta^1 = \frac{1}{2}S_1^2(\dot{p})$ :

$$\begin{array}{l} \log Pl = a^1 + \Delta^1 \\ \log p = a^1 \\ \log Pa = a^1 - \Delta^1 \end{array}$$

For small deviations in  $\dot{p}$  and  $\dot{q}$  the new and old variances of  $\dot{p}$  will approach equality,  $\Delta^1 = \Delta^0 = \Delta$ , not only absolutely but also relatively. This will normally hold accurately already for moderate deviations, because variances are relatively independent from their weighting (both have the same squared deviations of log-prices). For the variances in  $\dot{p}$  and  $\dot{q}$  both approaching zero (which makes old and new value shares to approach each other), we have asymptotic equations where  $\Delta$  clearly indicates 1 dose of bias measured from the central tine of the fork:

$$\begin{aligned} \log L &= a^0 + \Delta \\ \log l &= a^0 \\ \log Lh &= a^0 - \Delta \end{aligned}$$

$$\begin{aligned} \log Pl &= a^1 + \Delta \\ \log p &= a^1 \\ \log Pa &= a^1 - \Delta. \end{aligned}$$

From these equations, we see e.g. that  $\frac{1}{2}(\log L + \log Pa) = \log F = \frac{1}{2}(a^0 + a^1)$ , which equals e.g.  $\frac{1}{2}(\log l + \log p) = \log t = \frac{1}{2}(a^0 + a^1)$ . We calculate the geometric averages of between an index and its time antitheses (or equivalently the arithmetic averages of their log's) before equality  $\Delta^1 = \Delta^0 = \Delta$  and get:

$$\begin{aligned} \frac{1}{2}(\log L + \log Pa) &= \log F = \frac{1}{2}(a^0 + a^1) + \frac{1}{2}(\Delta^0 - \Delta^1) \rightarrow \frac{1}{2}(a^0 + a^1) \\ \frac{1}{2}(\log l + \log p) &= \log t = \frac{1}{2}(a^0 + a^1) \\ \frac{1}{2}(\log Lh + \log Pl) &= \log D = \frac{1}{2}(a^0 + a^1) - \frac{1}{2}(\Delta^0 - \Delta^1) \rightarrow \frac{1}{2}(a^0 + a^1) \end{aligned}$$

when  $\Delta^0 - \Delta^1$  and  $\frac{\Delta^0 - \Delta^1}{\Delta}$  approach zero. This happens if both  $\frac{1}{2}s^2(\dot{p}; w)$  and  $\frac{1}{2}s^2(\dot{q}; w)$  approach zero, for old and new value shares. This is a differential geometric argument, which becomes exact mathematics by using the shifts between the formulas, as is done in Appendix 1.

The point is that all the three rectified indices attain (for small *deviations* in  $\dot{p}$  and  $\dot{q}$ , the absolute *magnitude* of the log-changes is irrelevant) approximately the same value, and exactly so if the variances of  $\dot{p}$  and  $\dot{q}$  approach zero. The situation above describes what happens in the limit of almost the same proportional changes. This *common point*  $\frac{1}{2}(a^0 + a^1)$  defines the exact point of indices, the average point of these six indices on the log-scale, which can and should be declared as *unbiased*. Or more accurately, as *unbiased for almost proportional changes* of prices *and* of quantities. We have proved the following theorem (see Appendix 1): For all values of  $\log l = a^0$  and  $\log p = a^1$ , the indices ( $F, t, D$ ) approach each other and the constant  $\frac{1}{2}(a^0 + a^1)$  in the log-scale in the sense that their *Shifts* vanish when the variances of the log-changes in prices  $\Delta = \frac{1}{2}s^2(\dot{p}; \bar{w})$  and in quantities  $\Delta(q) = \frac{1}{2}s^2(\dot{q}; \bar{w})$  approach zero.

For large deviations, even these three precision formulas (or other asymptotically unbiased indices) start to scatter and one cannot say any more how the small biases start to distribute among them.

Here are only three TA-rectified new formulas, because e.g. the Törnqvist formula is attained by either rectifying of  $l$  or  $p$ . We do not present this as formula (or its simple algebra), because it is much better that the reader looks carefully the table 1 in Appendix 2 and clears to procedure to himself (or to herself).

For small variances in the log-changes of both prices and quantities, the three indices ( $F, t, D$ ) have approximately the same value, irrespective where the three forks situate ( $a^0, a^1$ ) and how they disperse  $\Delta$ . All other averages (than these TA-rectifications) over two members from different three-forks are *always biased* up or down. As proved in Vartia-Vartia (1984),  $\frac{1}{2}(\log L + \log Pl) = \frac{1}{2}(a^0 + a^1) + \Delta$  and therefore

both  $\sqrt{L * PL}$  and  $\frac{1}{2}(L + PL)$  have one dose of permanent bias up. Similarly, we get  $\frac{1}{2}(\log Lh + \log p) = \frac{1}{2}(a^0 + a^1) - \frac{1}{2}\Delta$  and thus  $\sqrt{Lh * p}$  has a half dose of permanent bias down, etc.

Only the three indices ( $F, t, D$ ) produced by TA-rectification are always unbiased and superlative of the ones we have discussed. Further research is needed to widen this important set of asymptotically unbiased index numbers, but these three indices have served (together with  $St$  and  $MV$ ) to show where the precision formulas lie and how near each other.

This ends our discussion and correction of the Fisher's partly experimental discussion of his concept of bias and unbiased index numbers. The main conclusion is that both Laspeyres  $L$  and Paasche  $Pa$  price indices are clearly (asymptotically and finitely) biased formulas, whose biases may be small only by accident.

As we have noted, the quantity side of data has much more variability than the price side. Therefore, we should routinely use the price indices as a starting point of the  $V = P * Q$  decomposition. For instance, start e.g. from  $t$  and use the pair  $(t, CoF(t))$  leading to

$$(15) \quad \frac{V^1}{V^0} = P * Q = t * CoF(t) .$$

This would give accurate results, instead of doing it the other way around,  $(P, Q) = (CoF(t), t)$ , where Törnqvist formula is first applied for quantities and then its co-factor  $CoF(t)$  is used as a price index. That would give useless results in 5S and poor results in 5N because the quantity indices vary too much. These FA-indices are totally useless in 5S because of the large variances  $s^2(\dot{q}; w)$  of the quantity log-changes  $\dot{q}$ . Their biases are really large real numbers, often outside  $\pm 5000$  doses of bias! Concretely, these large figures arise, because the FA-indices disperse so strongly. In many commodities, their quantity relatives approach infinity caused by division of nearly zero, and the index explodes, see Fig. 5 page 12.

On the other hand, the original six indices (calculated normally in the price-space) deviate only moderately from each other. Their graphs show that the old and new three-forks change abruptly places, especially in 2015-16, see Fig. 6 page 12.

In the set of normal commodities 5N, the price development is in total control, see Fig. 1 page 9.


## 9. Index number production for similar complete micro data

If we concentrate only on the precision formulas in 5S, the results are in good control. This will be most probably the case also in other similar detailed homogeneous price-quantity data sets, see Fig. 7 page 13.

The most important feature in our treatment was the division of data into two, as the union of normal and special commodities, say 5N and 5S. Simple like that! In 5N all the index calculations are quite easy and traditional and it helped us to define the separation between e.g. contingently biased, freakish and superlative formulas. From the superlative formulas, we mentioned only a few:  $F, t, St, D, MV, VW$ . Some others will be introduced in the next paper and their merits and dis-merits will be discussed. A concrete application of these formulas in measuring welfare change in consumer theory is Vartia (1983).

We shall analyze carefully these formulas and drop away (eliminate) those which do not react correctly to extreme changes (up or down) in prices and quantities, a less analyzed property discussed in Vartia (1976, 1976b). That kind of changes occur all the time in 5S and they become even more pronounced elements in 5S when new and disappearing commodities are added to it.

We hope that the future research will show, that the following picture describes the actual problem very accurately. If so, we can be indifferent between choosing among the indices in  $\{F, Stu\}$  in order to calculate the official price development in 5S and combine that with 5N using the same formula.



## 10. Conclusions

We analyze most typical index number formulas by the method of ‘Fisher forks’. We define mathematically the shift measure by which we may classify the index number formulas in to four classes: superlative, permanently biased, contingently biased and freakish. We visualize some time quite difficult mathematics by empirical founding’s and the graphs of them. In empirical analyze we use complete micro data including price, quantity and value information in all time periods.

Our data includes large deviations in quantity and/or price changes and naturally large variation in value shares. We show that only few index numbers can *tolerate* them. The situation is complicated in many ways, but logically arranged figures tell the story best. The concept of bias of the index number is very important issue of our study. It is well known, that there does not exist a simple numerical measure for bias or order of quality of an index number, because it depends on so many factors.

We have demonstrated, that Fisher’s theory, his Five-tined Fork,  $Pl > p > Pa \approx L > l > Lh$  is false because Laspeyres  $L$  and Paasche  $Pa$  are not even usually almost equal. Actually, index numbers fall in four groups, which are *Superlative*, *Permanently Biased*, *Contingently Biased* and *Freakish*. These are defined by the limit behavior of the function  $Shift(P, F)$  when changes (or deviations) in log-prices and log-quantities approach zero.

We have found that contingently biased indices, like simple base or observation period weighted indices (especially  $L$  and  $Pa$ ), get values here and there, but superlative indices go hand in hand. Choosing among the best formulas is more or less indifferent, because their values differ so little.

We analyze the data by classifying the commodities as normal ones, forming here the subset 5N, and special ones with exceptionally large relative changes in value. The latter formed the set of special commodities 5S in this paper, where individual values either increased 5-fold or declined to 1/5 of their old value. The subset 5S revealed simply and concretely, which index number formulas can be used in public information production. We are able to handle also the problems of new and disappearing commodities in a similar fashion.

In the forthcoming CPI, the number of commodities can be increased by factors 50-100 and also all the quantities will be observed for many subgroups of commodities, based on scanner data on all transactions on the most detailed commodity level. This "Big Data" will remove most of the problems in the former production system, where only some 1500 commodities out of roughly 100 000 actual ones were monitored and quantities were not observed at the same time. Other problems of index number construction possibly remaining are of minor importance.

The accuracy and reliability of CPI will rise to a higher level, without much additional costs.

Let us say farewell with thanks to the dear Laspeyres, which has been the prototype of decent index numbers. It has a number of good properties, but a single drawback: it is inaccurate.



## References

- Balk, Bert (1995): Axiomatic index number theory: A survey. *International Statistical Review*, 63, 1, 63-93.
- Cavallo, Alberto and Rigobon, Roberto (2016): The billion prices project: Using online prices for measurement and research, *Journal of Economic Perspectives*, 30, Nro 2, Spring 2016, 151-178.
- Diewert, Erwin (1976): Exact and superlative index numbers, *Journal of econometrics*, 4, 115-145.
- Diewert, Erwin (1978): Superlative index numbers and consistency in aggregation, *Econometrica*, 46, 883-900.
- Fisher, Irving (1922): **The making of index numbers**, Houghton Mifflin Company, Boston.
- Hansen, Bent and Lucas, Edward F. (1984): On the accuracy of index numbers, *Review of Income and Wealth*, 1984,1, 25-38.
- Montgomery, J.K. (1937): **The mathematical problem of the price index**, P.S. King et Son Ltd., London
- Pursiainen, Heikki (2005): **Consistent aggregation methods and index number theory**, Research Reports, Kansantaloustieteen laitoksen tutkimuksia (Research of the Department of Economics in Helsinki) No. 106:2005. Dissertationes Oeconomicae
- Sato, Kazuo (1976): The ideal log-change index number, *The Review of Economics and Statistics*, 58, 223-8.
- Stuvel, G. (1957): A new index number formula, *Econometrica*, 25, 123-131.
- Stuvel, G. (1989): **The Index-number problem and its solution**, Basinstoke and London, Macmillan.
- Suoperä, Antti and Vartia, Yrjö (2011): Analysis and synthesis of wage determination in heterogeneous gross sections, *HECER Discussion Papers No. 331, June 2011*.
- Törnqvist, Leo (1936): The Bank of Finland's consumption price index, *Bank of Finland Monthly Bulletin*, 10, 1-8.
- Törnqvist, Leo, Vartia, Pentti and Vartia, Yrjö (1985): How should the relative changes be calculated, *American Statistician*, 39, 1, 43-46.
- Vartia, Yrjö (1976): **Relative changes and index numbers**, Research Institute of the Finnish Economy ETLA, series A4 (dissertation in statistics).
- Vartia, Yrjö (1976b): Ideal log-change index numbers, *Scandinavian Journal of Statistics*, 3, 121-126.
- Vartia, Yrjö (1978): Fisher's five-tined fork and other quantum theories of index numbers, *In Theory and applications of economic indices* (Ed. W. Eichhorn, R. Henn, O. Opitz, R. W. Shephard), Physica-Verlag, Wurzburg.
- Vartia, Yrjö (1980): Compensated income in consumption-saving economies, Research reports No. 66, Research Institute of the Finnish Economy ETLA, 1-105.
- Vartia, Yrjö (1983): Efficient methods of measuring welfare change and compensated income in terms of ordinary demand functions, *Econometrica*, 51,79-98.
- Vartia, Yrjö (2008): On the aggregation of quadratic micro equations, *HECER Discussion Papers No. 248, December 2008*.
- Vartia, Yrjö (2010): Principles of defining index numbers and constructing index series, *HECER Discussion Papers No. 283, January 2010*.
- Vartia, Yrjö and Vartia, Pentti (1984): Descriptive index number theory and the Bank of Finland currency index, *Scandinavian Journal of Economics*, 86, 3, 352-364.
- Vartia, Yrjö and Weymark, John A. (1982): Four revealed preference tables, *Scandinavian Journal of Economics*, 83, 408-418.

## Appendix 1: Theorems on biasedness and unbiasedness

The relative weight bias depends on the difference of means  $\log p - \log l = \sum (w_i^1 - w_i^0) \dot{p}_i$  between the central lines of the new and old weighted forks. This can be written in several ways as an exact covariance between various variables. This rather surprising idea forms our first theorem.

**Theorem 1:**  $\log p - \log l = \sum (w_i^1 - w_i^0) \dot{p}_i = \sum \Delta w_i \Delta \log p_i$ .

This can be expressed as several different (but numerically equal) covariances. Thus, identically for any data:

$$\begin{aligned} \log p - \log l &= \sum \frac{1}{n} n \Delta w_i \Delta \log p_i = \text{cov} \left( n \Delta w, \dot{p}; \frac{1}{n} \right) && \text{even weights} \\ &= \sum w_i^0 \frac{\Delta w_i}{w_i^0} \Delta \log p_i = \text{cov} \left( \frac{\Delta w}{w^0}, \dot{p}; w^0 \right) && \text{old weights} \\ &= \sum \frac{1}{2} (w_i^0 + w_i^1) \left( \frac{\Delta w_i}{\bar{w}_i} \right) \Delta \log p_i = \text{cov} \left( \frac{\Delta w}{\bar{w}}, \dot{p}; \bar{w} \right) && \text{Törnqvist weights} \\ &= \sum L(w_i^0, w_i^1) \frac{\Delta w_i}{L(w_i^0, w_i^1)} \Delta \log p_i = S * \sum \frac{L(w_i^0, w_i^1)}{\sum L(w_i^0, w_i^1)} \Delta \log w_i * \Delta \log p_i = S * \text{cov}(\Delta \log w, \dot{p}; \hat{w}), \end{aligned}$$

where  $S = \sum L(w_i^0, w_i^1) \leq 1$  and  $\hat{w}_i = \frac{L(w_i^0, w_i^1)}{\sum L(w_i^0, w_i^1)}$  = weights of the Sato-Vartia index.

**Proof:** Covariance is defined as  $\text{cov}(x, y; w) = \sum w_i (x_i - \bar{x})(y_i - \bar{y}) = \sum w_i (x_i - \bar{x}) y_i$ , where the weights  $w$  are non-negative real numbers summing to unity and e.g.  $\bar{x} = \sum w_i x_i = w$ -weighted mean of the variable  $x$ . Thus,  $\sum w_i (x_i - \bar{x}) = 0$  is a characteristic property of any covariance. It implies  $\sum w_i (x_i - \bar{x})(y_i - \bar{y}) = \sum w_i (x_i - \bar{x}) y_i - \sum w_i (x_i - \bar{x}) * \bar{y} = \sum w_i (x_i - \bar{x}) y_i$ , i.e. only one of the variables  $x$  and  $y$  needs to be expressed in the deviation form!

In all the expression above this zero-condition of the first variable applies, as in  $\sum \frac{1}{2} (w_i^0 + w_i^1) \left( \frac{\Delta w_i}{\bar{w}_i} \right) = \sum \Delta w_i = 1 - 1 = 0$ . This makes them all covariances of the denoted variables and weights.

Next we prove the following intuitive results concerning the old and new variances  $s_0^2(\dot{p}) = s^2(\dot{p}, w^0)$  and  $s_1^2(\dot{p}) = s^2(\dot{p}, w^1)$  and their approximations by  $s^2(\dot{p}) = s^2(\dot{p}, \bar{w})$  with the Törnqvist weights. They all become equal, or more accurately, their ratios approach unity, when they themselves approach zero while log-changes in prices and quantities approach any proportional values. Their accurate mathematical expression is somewhat tricky:

**Theorem 2:** For all  $(\log P, \log Q) \in R^2: \dot{p} \rightarrow \log P * 1_n$  and  $\dot{q} \rightarrow \log Q * 1_n$  implies  $\frac{s_1^2(\dot{p}) - s_0^2(\dot{p})}{s^2(\dot{p})} \rightarrow 0$ .

The condition  $\dot{p} \rightarrow \log P * 1_n$  means that for all commodities the log-change  $\Delta \log p_i = \dot{p}_i$  approaches the same limit:  $\dot{p}_i = \Delta \log p_i \rightarrow \log P$ , or the log-deviation vector  $\dot{p} - \log P * 1_n$  approaches a zero vector. This refers to almost proportional changes APC.

**Proof:**  $s_1^2(\dot{p}) - s_0^2(\dot{p}) = \sum w_i^1 (\Delta \log p_i - \Delta \log p)^2 - \sum w_i^0 (\Delta \log p_i - \Delta \log l)^2 = \sum (w_i^1 - w_i^0) (\Delta \log p_i)^2 + (\Delta \log p)^2 - (\Delta \log l)^2 = \sum (w_i^1 - w_i^0) (\Delta \log p_i - \Delta \log t)^2$  by a straight calculation left to the reader as an exercise (expand, sum and note definitions of  $\Delta \log l$  and  $\Delta \log p$ ).  $|\sum (w_i^1 - w_i^0) (\Delta \log p_i - \Delta \log t)^2| = \left| \sum \bar{w}_i \left( \frac{\Delta w_i}{\bar{w}_i} \right) (\Delta \log p_i - \Delta \log t)^2 \right| \leq \sum \bar{w}_i \left| \frac{\Delta w_i}{\bar{w}_i} \right| (\Delta \log p_i - \Delta \log t)^2 \leq \sum \bar{w}_i \max \left| \frac{\Delta w_j}{\bar{w}_j} \right| (\Delta \log p_i - \Delta \log t)^2 = \sum \bar{w}_i (\Delta \log p_i - \Delta \log t)^2 * \max \left| \frac{\Delta w_j}{\bar{w}_j} \right| = s^2(\dot{p}) * \max \left| \frac{\Delta w_j}{\bar{w}_j} \right|$ . Thus  $\left| \frac{s_1^2(\dot{p}) - s_0^2(\dot{p})}{s^2(\dot{p})} \right| \leq \max \left| \frac{\Delta w_j}{\bar{w}_j} \right| = \max \left| \frac{L(w_j^0, w_j^1)}{\bar{w}_j} * \frac{\Delta w_j}{L(w_j^0, w_j^1)} \right| = \max \left| \frac{L(w_j^0, w_j^1)}{\bar{w}_j} * \Delta \log w_j \right| = \max \left| \frac{L(w_j^0, w_j^1)}{\bar{w}_j} (\dot{p}_j + \dot{q}_j) \right| \leq \max \left| \frac{L(w_j^0, w_j^1)}{\bar{w}_j} \right| * \max |(\dot{p}_j + \dot{q}_j)| \rightarrow 1 * 0 = 0$ , when all the deviations  $\dot{p}_i = \Delta \log p_i - \Delta \log t$  and  $\dot{q}_i = \Delta \log q_i - \Delta \log t_q$  approach zero (implying  $w_i^1 \rightarrow w_i^0$ ). This proves the theorem.

**Theorem 3:** For all  $(\log P, \log Q) \in R^2: \check{p} \rightarrow \log P * 1_n$  and  $\check{q} \rightarrow \log Q * 1_n$  implies  $\frac{s_1^2(\check{p}) + s_0^2(\check{p})}{2s^2(\check{p})} \rightarrow 1$ .

**Proof:**  $\frac{1}{2}(s_1^2(\check{p}) + s_0^2(\check{p})) = \frac{1}{2}\sum w_i^1(\Delta \log p_i - \Delta \log p)^2 + \frac{1}{2}\sum w_i^0(\Delta \log p_i - \Delta \log l)^2$   
 $= \frac{1}{2}\sum(w_i^1 + w_i^0)(\Delta \log p_i)^2 - (\Delta \log p)^2 - (\Delta \log l)^2 = \frac{1}{2}\sum(w_i^1 + w_i^0)(\Delta \log p_i - \Delta \log t)^2 + (\Delta \log t)^2 - (\Delta \log p)^2 -$   
 $(\Delta \log l)^2 = s^2(\check{p}) - \frac{1}{4}(\log p - \log l)^2 = s^2(\check{p}) - \frac{1}{4}\left(\text{cov}\left(\frac{\Delta w}{\bar{w}}, \check{p}; \bar{w}\right)\right)^2$ . Thus  $\left|\frac{s_1^2(\check{p}) + s_0^2(\check{p})}{2s^2(\check{p})}\right| = \left|1 - \frac{1}{4s^2(\check{p})}\left(\text{cov}\left(\frac{\Delta w}{\bar{w}}, \check{p}; \bar{w}\right)\right)^2\right| \leq 1 + \left|\frac{1}{4}\left(s^2\left(\frac{\Delta w}{\bar{w}}; \bar{w}\right)r^2\left(\frac{\Delta w}{\bar{w}}, \check{p}; \bar{w}\right)\right)\right| \leq \frac{1}{4}s^2\left(\frac{\Delta w}{\bar{w}}; \bar{w}\right) \rightarrow 0$  when all the deviations  $\check{p}_i = \Delta \log p_i - \Delta \log t$  and  $\check{q}_i = \Delta \log q_i - \Delta \log t_q$  approach zero (implying  $w_i^1 \rightarrow w_i^0$  for all  $i$ ). This proves the theorem.

**Theorem 4:** For all  $(\log P, \log Q) \in R^2: \check{p} \rightarrow \log P * 1_n$  and  $\check{q} \rightarrow \log Q * 1_n$  implies

$$\frac{s_1^2(\check{p})}{s^2(\check{p})} \rightarrow 1 \text{ and } \frac{s_0^2(\check{p})}{s^2(\check{p})} \rightarrow 1.$$

**Proof:** Insert identity  $s_1^2(\check{p}) = \frac{1}{2}(s_1^2(\check{p}) + s_0^2(\check{p})) + \frac{1}{2}(s_1^2(\check{p}) - s_0^2(\check{p}))$  into  $\frac{s_1^2(\check{p})}{s^2(\check{p})} = \frac{s_1^2(\check{p}) + s_0^2(\check{p})}{2s^2(\check{p})} + \frac{s_1^2(\check{p}) - s_0^2(\check{p})}{2s^2(\check{p})}$ .  
 By theorems 2 and 3, this approaches  $1+0 = 1$  for APC, which proves  $\frac{s_1^2(\check{p})}{s^2(\check{p})} \rightarrow 1$ . Similarly  $\frac{s_0^2(\check{p})}{s^2(\check{p})} \rightarrow 1$ .

Now we are ready to state and prove the main result that Törnqvist  $t$  and Fisher  $F$  approximate each other quadratically for almost proportional changes and thus for all small changes. The proof is short and elegant:

**Theorem 5:** For all  $(\log P, \log Q) \in R^2: \check{p} \rightarrow \log P * 1_n$  and  $\check{q} \rightarrow \log Q * 1_n$  the limes  $\lim_{APC} Shift(t, F)$  exists and equals zero. Thus, Törnqvist  $t$  is superlative for almost proportional changes. Therefore, it is also unbiased for small changes or  $\lim_{SC} Shift(t, F) = 0$ .

**Proof:** By the properties of moment means

$$A \quad \log Pa - \log p = -\frac{1}{2}s_1^2(\check{p}) + o_2 \text{ and } \log L - \log l = \frac{1}{2}s_0^2(\check{p}) + o_2,$$

where  $o_2 = o_2(\check{p}, \check{q})$  means any expression, which goes to zero faster than the variance (or sum of squares) of  $\check{p}$  when both  $(\check{p}, \check{q})$  approach zero, i.e.  $s^2(\check{p}) \rightarrow 0$  and  $s^2(\check{q}) \rightarrow 0$ . Thus  $o_2/s^2(\check{p}) \rightarrow 0$  when the denominator and  $s^2(\check{q})$  approach zero. Summing equations A, and dividing by two gives

$$\log F - \log t = -\frac{1}{4}(s_1^2(\check{p}) - s_0^2(\check{p})) + o_2. Shift(t, F) = \frac{2 \log\left(\frac{t}{F}\right)}{s^2(\check{p}, \bar{w})} = \frac{1}{2} \frac{(s_1^2(\check{p}) - s_0^2(\check{p}))}{s^2(\check{p}, \bar{w})} + \frac{o_2}{s^2(\check{p}, \bar{w})}. \text{ Now by}$$

Theorem 1,  $\lim_{APC} Shift(t, F) = 0 + 0 = 0$ .

**Theorem 6:** The moment means  $M^{(\alpha)}(p^{1/0}, \bar{w}) = \left(\sum \frac{1}{2}(w_i^0 + w_i^1)\left(\frac{p_i^1}{p_i^0}\right)^\alpha\right)^{1/\alpha}$  with Törnqvist weights are permanently biased upwards for all  $\alpha > 0$  and downwards for  $\alpha < 0$ . This is an infinite set (continuum) of permanently biased indices. The permanent bias, which equals  $\alpha$ , is small only when it is nearly zero (the superlative Törnqvist index case). This implies that the Törnqvist weights (average value shares) are correct weights only for the geometric index (= Törnqvist index).

**Proof:** Apply (4) for Törnqvist weights and get  $\log \frac{M^{(\alpha)}(p^{1/0}, \bar{w})}{t} = \alpha s^2(\check{p}, \bar{w}) + o_2$  and  $Shift(M^{(\alpha)}(p^{1/0}, \bar{w}), t) = Shift(M^{(\alpha)}(p^{1/0}, \bar{w}), F) + Shift(F, t) = \alpha + Shift(F, t) + o_2/s^2(\check{p}, \bar{w}) \rightarrow \alpha$ , by theorem 5 when both  $(\check{p}, \check{q})$  approach zero. Thus, its limit exists and equals a constant  $\alpha$ , which is the value of the permanent bias. This proves the theorem.

**Theorem 7:**  $\log\left(\frac{p}{t}\right) = \text{cov}(\check{p}, \check{p}; \bar{w}) + o_2 = \text{cov}(\check{p}, \check{q}; \bar{w}) + s^2(\check{p}, \bar{w}) + o_2$  and therefore, e.g.

$Shift(p, t) \rightarrow Shift(p, F) \rightarrow \frac{cov(\dot{v}, \dot{p}; \bar{w})}{s^2(\dot{p}, \bar{w})}$  when deviations in price and quantity log-changes approach zero or APC.

**Proof:** Recall the basic properties of cosh and sinh:

$$\cosh(z) = \frac{1}{2}(\exp(z) + \exp(-z)) = 1 + \frac{1}{2!}z^2 + \frac{1}{4!}z^4 + \dots \text{ and}$$

$$\sinh(z) = \frac{1}{2}(\exp(z) - \exp(-z)) = z + \frac{1}{3!}z^3 + \frac{1}{5!}z^5 + \dots$$

First  $\bar{w}/\sqrt{w^0 w^1} = \frac{1}{2}(\sqrt{\frac{w^1}{w^0}} + \sqrt{\frac{w^0}{w^1}}) = \frac{1}{2}(\exp(\frac{1}{2}\log(\frac{w^1}{w^0})) + \exp(-\frac{1}{2}\log(\frac{w^1}{w^0}))) = \frac{1}{2}(\exp(z) + \exp(-z))$

$$= \cosh(z) \text{ for } z = \frac{1}{2}\log\left(\frac{w^1}{w^0}\right) = \frac{1}{2}\Delta\log w. \text{ Thus } \frac{\bar{w}}{\sqrt{w^0 w^1}} = 1 + \frac{1}{8}\log\left(\frac{w^1}{w^0}\right)^2 + \dots = 1 + \frac{1}{8}\ddot{w}^2 + o_2.$$

Similarly  $\frac{L(w^0, w^1)}{\sqrt{w^0 w^1}} = L\left(\sqrt{\frac{w^1}{w^0}}, \sqrt{\frac{w^0}{w^1}}\right) = L(\exp(\frac{1}{2}\log(\frac{w^1}{w^0})), \exp(-\frac{1}{2}\log(\frac{w^1}{w^0}))) = L(\exp(z), \exp(-z)) = \sinh(z)/z. \text{ Thus}$

$$\frac{L(w^0, w^1)}{\sqrt{w^0 w^1}} = 1 + \frac{1}{3!}z^2 + \dots = 1 + \frac{1}{24}\ddot{w}^2 + o_2 \text{ and } \frac{L(w^0, w^1)}{\bar{w}} = \frac{L(w^0, w^1)\sqrt{w^0 w^1}}{\sqrt{w^0 w^1}\bar{w}} = \frac{L(w^0, w^1)}{\sqrt{w^0 w^1}} / \frac{\bar{w}}{\sqrt{w^0 w^1}}.$$

Therefore  $\frac{L(w^0, w^1)}{\bar{w}} = 1 + \frac{1}{24}\ddot{w}^2 - \frac{1}{8}\ddot{w}^2 + \dots = 1 - \frac{1}{12}\ddot{w}^2 + \dots$  and by theorem 1  $\log p - \log l = \sum \frac{1}{2}(w_i^0 + w_i^1) \left(\frac{\Delta w_i}{w_i}\right) \dot{p}_i = \sum \frac{1}{2}(w_i^0 + w_i^1) \left(\frac{\Delta w_i}{L(w_i^0, w_i^1)}\right) \frac{L(w_i^0, w_i^1)}{\bar{w}_i} \dot{p}_i = \sum \frac{1}{2}(w_i^0 + w_i^1) \Delta \log w_i (1 + \frac{1}{12}\ddot{w}_i^2) \dot{p}_i + o_2 = \sum \frac{1}{2}(w_i^0 + w_i^1) \Delta \log v_i \dot{p}_i + o_2 = cov(\dot{v}, \dot{p}; \bar{w}) + o_2.$  Insert this in  $Shift(p, t) = \frac{\log(\frac{p}{t})}{\frac{1}{2}s^2(\dot{p}, \bar{w})} = \frac{\frac{1}{2}\log(\frac{p}{t})}{\frac{1}{2}s^2(\dot{p}, \bar{w})} = \frac{cov(\dot{v}, \dot{p}; \bar{w})}{s^2(\dot{p}, \bar{w})} + \frac{o_2}{s^2(\dot{p}, \bar{w})}.$  Now let deviations in price and quantity log-changes approach zero without altering their essential mutual dependencies, such as correlation and the ratio of deviations. Then  $Shift(p, t) \rightarrow \frac{cov(\dot{v}, \dot{p}; \bar{w})}{s^2(\dot{p}, \bar{w})}$ , which is not a unique real number, but a contingent data-dependent function. Or alter them to get some *other* limit. This proves that a unique limit does not exist and  $p$  is contingently biased in respect to  $t$  or equivalently in respect to Fisher. This proves the theorem for APC and thus for SC.

**Theorem 8:**  $\log\left(\frac{Pa}{L}\right) = cov(\dot{p}, \dot{q}; \bar{w}) + o_2$  and therefore, e.g.  $Shift(Pa, F) \rightarrow \frac{cov(\dot{p}, \dot{q}; \bar{w})}{s^2(\dot{p}, \bar{w})}$  when deviations in price and quantity log-changes approach zero or APC.

**Proof:** Subtracting equations A gives by theorems 2 and 6:  $\log Pa - \log L = \log p - \log l - \frac{1}{2}(s_1^2(\dot{p}) + s_0^2(\dot{p})) + o_2 = cov(\dot{v}, \dot{p}; \bar{w}) - s^2(\dot{p}, \bar{w}) + o_2 = cov(\dot{p}, \dot{q}; \bar{w}) + o_2.$  The  $Shift(Pa, F)$  and its limit behavior follows just like in theorem 7.

**Theorem 9:** Drobish  $Dr = \frac{1}{2}(L + Pa)$  is superlative for APC and thus for SC. It has a very appealing basket interpretation  $Dr = \frac{p^1 \cdot (q^0 + q^1/L(q))}{p^0 \cdot (q^0 + q^1/L(q))}$  in contrast to Fisher. It is an example of a superlative index fulfilling neither Time Reversal nor Quantity Reversal Tests.

**Proof:**  $\frac{Dr}{F} = \frac{1}{2}\left(\frac{L}{F} + \frac{Pa}{F}\right) = \frac{1}{2}\left(\sqrt{\frac{L}{Pa}} + \sqrt{\frac{Pa}{L}}\right) = \frac{1}{2}\left(\exp\left(\frac{1}{2}\log\frac{L}{Pa}\right) + \exp\left(-\frac{1}{2}\log\frac{L}{Pa}\right)\right) = \frac{1}{2}(\exp(z) + \exp(-z))$

$$= \cosh(z) = 1 + \frac{1}{2!}z^2 + \frac{1}{4!}z^4 + \dots = 1 + \frac{1}{8}\left(\log\frac{L}{Pa}\right)^2 + o_2 \text{ as in theorem 7.}$$

$$\log\left(\frac{Dr}{F}\right) = \log\left(1 + \frac{1}{8}\left(\log\frac{L}{Pa}\right)^2 + o_2\right) = \frac{1}{8}\left(\log\frac{L}{Pa}\right)^2 + o_2 = \frac{1}{8}cov(\dot{p}, \dot{q})^2 + o_2 \text{ by theorem 8.}$$

$$Shift(Dr, F) = \frac{2\log\left(\frac{Dr}{F}\right)}{s^2(\dot{p}, \bar{w})} = \frac{1}{4}\frac{cov(\dot{p}, \dot{q}; \bar{w})^2 + o_2}{s^2(\dot{p}, \bar{w})} = \frac{1}{4}\frac{(s(\dot{p})s(\dot{q})r(\dot{p}, \dot{q}))^2 + o_2}{s^2(\dot{p})} = \frac{1}{4}s^2(\dot{q})r(\dot{p}, \dot{q})^2 + \frac{o_2}{s^2(\dot{p})}.$$

This approaches zero when for all  $(\log P, \log Q) \in \mathbb{R}^2: \dot{p} \rightarrow \log P * 1_n$  and  $\dot{q} \rightarrow \log Q * 1_n$  or when  $s^2(\dot{p})$  and  $s^2(\dot{q})$  both go to zero. We have proved that  $\lim_{APC} Bias(Dr, F)$  exists and equals zero. This means, that Drobish is unbiased for APC and thus for SC.

**Theorem 10:** Similarly constructed logarithmic mean  $L(L, Pa)$  of Laspeyres and Paasche, which lies between Dr and F (and is thus even nearer to F than Dr) is superlative for APC and thus for SC. It does not have a basket interpretation and does not fulfill Time Reversal or Quantity Reversal Tests.

**Proof:** Because  $Dr \geq L(L, Pa) \geq F$  also  $\log\left(\frac{Dr}{F}\right) \geq \log\left(\frac{L(L, Pa)}{F}\right) \geq 0$  and  $Shift(Dr, F) \geq Shift(L(L, Pa), F) \geq 0$ . Therefore, by the previous theorem, both shifts approach zero when  $s^2(\dot{p})$  and  $s^2(\dot{q})$  both approach zero.

**Theorem 11:** Any evenly weighted moment mean  $M^{(\alpha)}(L, Pa)$  or  $M^{(\alpha)}(l, p)$  with parameter  $\alpha \in [0, 1]$  is superlative for APC and thus for SC. For the parameter  $\alpha$  smaller than 1 (= the arithmetic mean), the moment mean is nearer to F and t. This produces an infinite number of superlative indices.

**Proof:**  $M^{(\alpha)}(L, Pa)$  proved like in Theorem 10 and  $M^{(\alpha)}(l, p)$  like in theorem 9.

**Theorem 12:** ‘‘Vartia-Walsh’’  $\log VW = \sum \sqrt{w_i^0 w_i^1} \Delta \log p_i$  is superlative for SC but not for APC.

**Proof:** From theorem 6

$\frac{\sqrt{w^0 w^1}}{\bar{w}} = 1 - \frac{1}{8} \ddot{w}^2 + o_2$  and  $\frac{1}{8} \ddot{w}^2 = 1 - \frac{\sqrt{w^0 w^1}}{\bar{w}} + o_2$ . For simplicity, we drop the commodity sub-indices.

$$\begin{aligned} \log(t) - \log VW &= \sum \bar{w} \Delta \log p - \sum \sqrt{w^0 w^1} \Delta \log p \\ &= \sum (\bar{w} - \sqrt{w^0 w^1}) \Delta \log p = \sum \bar{w} \left(1 - \frac{\sqrt{w^0 w^1}}{\bar{w}}\right) \Delta \log p = -\frac{1}{8} \sum \bar{w} \ddot{w}^2 \Delta \log p + o_2 \end{aligned}$$

Here the dotted variables are log-deviations from mean, not mere log-changes. This difference causes first difficulties.  $Shift(VW, t) = \frac{-\frac{1}{8} \sum \bar{w} \ddot{w}^2 \Delta \log p + o_2}{\frac{1}{2} s^2(\dot{p})}$  and  $-4Shift(VW, t) = \frac{\sum \bar{w} \ddot{w}^2 \Delta \log p + o_2}{s^2(\dot{p})}$ .

$\sum \bar{w} \ddot{w}^2 = \sum \bar{w} (\ddot{v} - \log \frac{v^1}{v^0})^2 = \sum \bar{w} \dot{v}^2 + o_2 = var(\ddot{v}; \bar{w}) + o_2 = \sum \bar{w} (\ddot{p}^2 + 2\ddot{p}\ddot{q} + \ddot{q}^2) + o_2 = var(\ddot{p}) + 2cov(\ddot{p}, \ddot{q}) + var(\ddot{q}) + o_2$ . Here we could equally well have single dots instead of double dots, because e.g.  $var(\ddot{p}) = var(\dot{p})$ . By triangle inequality  $\left| \frac{\sum \bar{w} \ddot{w}^2 \Delta \log p + o_2}{s^2(\dot{p})} \right| \leq \frac{\sum \bar{w} \ddot{w}^2 |\Delta \log p| + |o_2|}{s^2(\dot{p})} \leq \frac{(\sum \bar{w} \ddot{w}^2) \max |\Delta \log p| + |o_2|}{s^2(\dot{p})} \leq \left(1 + 2 \frac{s(\dot{q})}{s(\dot{p})} + \frac{s^2(\dot{q})}{s^2(\dot{p})}\right) \max |\Delta \log p| + \frac{o_2}{s^2(\dot{p})}$ . Suppose that that the log-changes approach zero so that  $\frac{s^2(\dot{q})}{s^2(\dot{p})} \leq K$ , say  $\leq 400$ . Then  $\left| \frac{\sum \bar{w} \ddot{w}^2 \Delta \log p + o_2}{s^2(\dot{p})} \right| \leq (1 + 2\sqrt{K} + K) \max |\Delta \log p| + \frac{o_2}{s^2(\dot{p})}$ , which approaches zero when all changes of log-prices and log-quantities approach zero. Because this holds for all choices of K, it does not restrict the limit processes in any way and  $Lim_{SC} Shift(VW, t) = Lim_{SC} Shift(VW, F) = 0$ .

**Corollary:** ‘‘Vartia-Walsh’’  $\log VW = \sum \sqrt{w_i^0 w_i^1} \Delta \log p_i$  is probably the simplest superlative index for SC which is consistent in aggregation CA. Its weights have many interesting expressions, but sum at most to unity.

**Theorem 13:** Montgomery-Vartia index  $\log MV = \sum \hat{w}_i \Delta \log p_i$ ,  $\hat{w}_i = L(v_i^0, v_i^1)/L(v^0, v^1)$  is superlative for SC but not for APC.

**Proof:** By similar argumentation as above  $Shift(MV, t) = \frac{-\frac{1}{24} \sum \bar{w} \ddot{w}^2 \Delta \log p + o_2}{\frac{1}{2} s^2(\dot{p})}$  from which the claim follows similarly. It is usually nearer to Törnqvist than ‘‘Vartia-Walsh’’.

**Theorem 14:** Consider any two superlative indices  $P_1$  and  $P_2$ . Any value between them may be produced by some superlative index. Especially, their weighted mean (or convex combination)  $P(\theta) = \theta P_1 + (1-\theta)P_2$ , where  $0 \leq \theta \leq 1$  is such a superlative index.

**Proof:** Because  $0 \leq |\log(P(\theta), F)| \leq \max |\log(P_i, F)|$ , both the corresponding shifts approach zero by assumption of superlativity of  $P_1$  and  $P_2$ .

**Theorem 15:** Consider geometric means between any two old and new weighted moment mean indices (expressed in log-form)  $\log P_0(\alpha) = \frac{1}{\alpha} \log \sum w_i^0 \left(\frac{p_i^1}{p_i^0}\right)^\alpha$  and  $\log P_1(\alpha) = \frac{1}{\alpha} \log \sum w_i^1 \left(\frac{p_i^1}{p_i^0}\right)^\alpha$ . Especially, combine  $\log L = \log P_0(1)$  with  $\log P_1(-1 + \delta)$ , which produce  $\sqrt{P_0(1)P_1(-1 + \delta)}$ . For  $\delta = 0$  this is the *superlative* index  $F$ , because  $P_1(-1) = TA(P_0(1))$ . For all other values of  $\delta$  it has a *permanent bias*, which is upwards if  $\delta > 0$  and downwards if  $\delta < 0$ . All this is expressed shortly as  $Shift_{APC}(\sqrt{P_0(1)P_1(-1 + \delta)}, F) = \frac{\delta}{2}$ .

This is a special case  $\alpha = 1$  of the next theorem.

**Theorem 16:** Consider geometric means between any two old and new weighted moment mean indices (expressed in log-form)  $\log P_0(\alpha) = \frac{1}{\alpha} \log \sum w_i^0 \left(\frac{p_i^1}{p_i^0}\right)^\alpha$  and  $\log P_1(\alpha) = \frac{1}{\alpha} \log \sum w_i^1 \left(\frac{p_i^1}{p_i^0}\right)^\alpha$ . Especially, combine  $\log P_0(\alpha)$  with  $\log P_1(-\alpha + \delta)$ , which produce  $\sqrt{P_0(\alpha)P_1(-\alpha + \delta)}$ . For  $\delta = 0$  this is a *superlative* index, because  $P_1(-\alpha) = TA(P_0(\alpha))$ . For all other values of  $\delta$  it has a *permanent bias*, which is upwards if  $\delta > 0$  and downwards if  $\delta < 0$ . All this is expressed shortly as  $Shift_{APC}(\sqrt{P_0(\alpha)P_1(-\alpha + \delta)}, F) = \frac{\delta}{2}$ .

**Proof:** We have  $\log \frac{P_0(\alpha)}{P_0(0)} = \frac{\alpha}{2} s_0^2(\dot{p}) + o_2$  and  $\log \frac{P_1(-\alpha + \delta)}{P_1(0)} = \frac{-\alpha + \delta}{2} s_1^2(\dot{p}) + o_2$ . Summing and dividing by two gives  $\log \sqrt{P_0(\alpha)P_1(-\alpha + \delta)} - \log t = \frac{\alpha}{4} s_0^2(\dot{p}) + \frac{-\alpha + \delta}{4} s_1^2(\dot{p}) + o_2 = \frac{\alpha}{4} (s_0^2(\dot{p}) - s_1^2(\dot{p})) + \frac{\delta}{4} s_1^2(\dot{p}) + o_2$ . By theorems 2, 4 and 5:  $Shift_{APC}(\sqrt{P_0(\alpha)P_1(-\alpha + \delta)}, F) = \frac{\delta}{2}$ .

**Corollary:** All indices of the form  $\sqrt{P_0(\alpha)P_1(\beta)}$  are *permanently biased*, unless if  $\beta = -\alpha$ , when they are *superlative*. In concrete applications parameters are usually small, roughly  $\alpha, \beta \in [-2, 2]$  or even  $[-1, 1]$ .

**Theorem 17:** Evenly weighted geometric mean of price relatives  $Ge$  and similar geometric mean with arbitrary constant weights  $Ge(c)$  are freakish for small changes and have a serious contingent bias.

**Proof:** We have to show only that  $Shift(Ge, t)$  approaches  $\pm\infty$  for some arguments  $\dot{p}, \dot{q}$  approaching zero. We have  $\log Ge = \sum \frac{1}{n} \Delta \log p_i$ ,  $\log(Ge/t) = \sum (\frac{1}{n} - \bar{w}_i) \Delta \log p_i$  and  $s^2(\dot{p}) = \sum \bar{w}_i (\Delta \log p_i)^2 - (\sum \bar{w}_i \Delta \log p_i)^2$ . Suppose that only the components of one commodity  $i$  differ from zero. These imply  $Shift(Ge, To) = \frac{2(\frac{1}{n} - \bar{w}_i) \Delta \log p_i}{s^2(\dot{p})} = \frac{2(\frac{1}{n} - \bar{w}_i) \Delta \log p_i}{\bar{w}_i (\Delta \log p_i)^2 - (\bar{w}_i \Delta \log p_i)^2} = \frac{2(\frac{1}{n} - \bar{w}_i)}{\bar{w}_i (1 - \bar{w}_i) \Delta \log p_i} \rightarrow \pm\infty$  (if  $\frac{1}{n} \neq \bar{w}_i > 0$ ) when  $\Delta \log p_i \rightarrow 0$ . Because the even weight differs from the Törnqvist weight, the shift of the evenly weighted geometric mean  $Ge$  explodes. It has a serious contingent bias. The result for  $\log Ge(c) = \sum c_i \Delta \log p_i$ ,  $\sum c_i = 1$ ,  $c_i \geq 0$  is proved similarly.

## Appendix 2: A 10-fork of six basic formulas and their FA's.

All these  $12 - 2 = 10$  indices ( $L$  and  $Pa$  appear twice in 12) can be estimated (with increasing accuracy for progressively smaller changes) using only four parameters:

$$m = \log t, \quad d = \log\left(\frac{p}{l}\right) \cong \text{cov}(\dot{v}, \dot{p}; \bar{w}), \quad \Delta(p) = \text{cov}(\dot{p}, \dot{p}; \bar{w})/2, \quad \Delta(q) = \text{cov}(\dot{q}, \dot{q}; \bar{w})/2.$$

We have recursively  $\log p = m + d/2$ ,  $\log l = m - d/2$ ,  $\log Pl = \log p + \Delta(p)$ ,  $\log Pa = \log p - \Delta(p)$ ,  $\log L = \log l + \Delta(p)$ ,  $\log Lh = \log l - \Delta(p)$ .

FA's differ from each other by  $\Delta(q)$ :  $\log FA(L) = \log Pa$ ,  $\log FA(Pa) = \log L$ ,  $\log FA(l) = \log Pa + \Delta(q)$ ,  $\log FA(p) = \log L - \Delta(q)$ ,  $\log FA(Lh) = \log Pa + 2\Delta(q)$ ,  $\log FA(Pl) = \log L - 2\Delta(q)$ .

All these indices are clearly biased, more or less depending on the situation. They all can be corrected for bias by TA-rectification, which implies that none of them should be ever used (in case of complete micro data). This is the starting point of our new "quantum theory of index numbers" for six base indices and their FA's, which replaces the false "Fisher's Five-tined Fork". It must be completed by including "the center of gravity"  $m = \log t$  and other new points corresponding to derivative indices. It is a rather complicated set of points and gets different forms for different values of its four parameters ( $m, d, \Delta(p), \Delta(q)$ ). It reduces to Fisher's Five-tined Fork for the two-parametric case ( $m, d = 2\Delta(p) = 2\Delta(q)$ ).

## Appendix 3: TA and FA for the basic six formulas

Table 1: Time Antithesis formulas TA for the basic six formulas.

(1)	(2)	(3)	(4)	(5)
	$p^{1/0}$	$p^{0/1}$	$1/p^{0/1}$	
Symbol of the formula	$f\left(\begin{smallmatrix} p^1 & q^1 \\ p^0 & q^0 \end{smallmatrix}\right)$	$f\left(\begin{smallmatrix} p^0 & q^0 \\ p^1 & q^1 \end{smallmatrix}\right)$	$1/f\left(\begin{smallmatrix} p^0 & q^0 \\ p^1 & q^1 \end{smallmatrix}\right)$	Symbol of the TA formula
$L$	$\frac{p^1 \cdot q^0}{p^0 \cdot q^0}$	$\frac{p^0 \cdot q^1}{p^1 \cdot q^1}$	$\frac{p^1 \cdot q^1}{p^0 \cdot q^1}$	$Pa = TA(L)$
$l$	$\prod \left(\frac{p_i^1}{p_i^0}\right)^{w_i^0}$	$\prod \left(\frac{p_i^0}{p_i^1}\right)^{w_i^1}$	$\prod \left(\frac{p_i^1}{p_i^0}\right)^{w_i^1}$	$p = TA(l)$
$Lh$	$1/\sum w_i^0 p_i^{0/1}$	$1/\sum w_i^1 p_i^{1/0}$	$\sum w_i^1 p_i^{1/0}$	$Pl = TA(Lh)$
$Pl$	$\sum w_i^1 p_i^{1/0}$	$\sum w_i^0 p_i^{0/1}$	$1/\sum w_i^0 p_i^{0/1}$	$Lh = TA(Pl)$
$p$	$\prod \left(\frac{p_i^1}{p_i^0}\right)^{w_i^1}$	$\prod \left(\frac{p_i^0}{p_i^1}\right)^{w_i^0}$	$\prod \left(\frac{p_i^1}{p_i^0}\right)^{w_i^0}$	$l = TA(p)$
$Pa$	$\frac{p^1 \cdot q^1}{p^0 \cdot q^1}$	$\frac{p^0 \cdot q^0}{p^1 \cdot q^0}$	$\frac{p^1 \cdot q^0}{p^0 \cdot q^0}$	$L = TA(Pa)$

Table 2: Factor Antithesis formulas *FA* for the basic six formulas.

(1)	(2)	(3)	(4)	(5)
	$p^{1/0}$	$P^{1/0}(q)$	$CoF(P^{1/0}(q))$	
Symbol of the formula	$f\left(\frac{p^1 q^1}{p^0 q^0}\right)$	$f\left(\frac{q^1 p^1}{q^0 p^0}\right)$	$V^{1/0}/f\left(\frac{q^1 p^1}{q^0 p^0}\right)$	Symbol of the FA formula
<i>L</i>	$\frac{p^1 \cdot q^0}{p^0 \cdot q^0}$	$\frac{q^1 \cdot p^0}{q^0 \cdot p^0}$	$\frac{p^1 \cdot q^1}{p^0 \cdot q^1}$	<i>Pa</i> = <i>FA(L)</i>
<i>l</i>	$\prod \left(\frac{p_i^1}{p_i^0}\right)^{w_i^0}$	$\prod \left(\frac{q_i^1}{q_i^0}\right)^{w_i^0}$	$V^{1/0} / \prod \left(\frac{q_i^1}{q_i^0}\right)^{w_i^0}$	<i>FA(l)</i>
<i>Lh</i>	$1 / \sum w_i^0 p_i^{0/1}$	$1 / \sum w_i^0 q_i^{0/1}$	$V^{1/0} / \sum w_i^0 q_i^{0/1}$	<i>FA(Lh)</i>
<i>Pl</i>	$\sum w_i^1 p_i^{1/0}$	$\sum w_i^1 q_i^{1/0}$	$V^{1/0} / \sum w_i^1 q_i^{1/0}$	<i>FA(Pl)</i>
<i>p</i>	$\prod \left(\frac{p_i^1}{p_i^0}\right)^{w_i^1}$	$\prod \left(\frac{q_i^1}{q_i^0}\right)^{w_i^1}$	$V^{1/0} / \prod \left(\frac{q_i^1}{q_i^0}\right)^{w_i^1}$	<i>FA(p)</i>
<i>Pa</i>	$\frac{p^1 \cdot q^1}{p^0 \cdot q^1}$	$\frac{q^1 \cdot p^1}{q^0 \cdot p^1}$	$\frac{p^1 \cdot q^0}{p^0 \cdot q^0}$	<i>L</i> = <i>FA(Pa)</i>